

The following manuscript was accepted for publication in Pharmaceutical Sciences. It is assigned to an issue after technical editing, formatting for publication and author proofing

Citation: Dehmoradkhani KR, Jafari A, Shokoohinia Y, Emami SA, Mojarrab M. Flavonoids from the aerial parts of *Artemisia biennis* Willd, Pharm. Sci. 2019, doi:10.15171/PS.2019.33

Short Communication

Flavonoids from the aerial parts of *Artemisia biennis* Willd.

Running title: Flavonoids from *Artemisia biennis*

Kazhal Rostami Dehmoradkhani ^a, Azizollah Jafari ^b, Yalda Shokoohinia ^c, Seyed Ahmad Emami ^d, Mahdi Mojarrab ^{*c}

^a Department of Chemistry, Faculty of sciences, University of Yasouj, Yasouj, Iran.

^b Department of Biology, Faculty of sciences, University of Yasouj, Yasouj, Iran.

^c Pharmaceutical Sciences Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran

^d Department of Pharmacognosy, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

*Corresponding author:

Mahdi Mojarrab

Associate professor of Pharmacognosy

Department of Pharmacognosy and Pharmaceutical Biotechnology

School of Pharmacy

Kermanshah University of Medical Sciences

Kermanshah

Iran

Tel: +98-833-4276480

Fax: +98-833-4276483

P.O. Box 67145- 1673

email addresses: mahdi.mojarrab@gmail.com; mmojarrab@kums.ac.ir

Abstract

Background: The current phytochemical study was carried out on a fraction of dried polar extract of aerial parts of *Artemisia biennis* Willd. which was previously reported to decrease the viability of the rat pheochromocytoma cell line PC12 in cell-based antioxidant assays. **Methods:** A combination of solid phase extraction (SPE) and high pressure liquid chromatography (HPLC) of the hydroethanolic extract was used to purify the compounds. Structures of the isolated compounds were elucidated by spectroscopic means, including ¹HNMR and MS analyses. **Results:** Three isolated and identified flavonoids in this study were luteolin, kaempferol and apigenin. **Conclusion:** The cytotoxic potential of flavone aglycones, as the major components of selected fraction in the hydroethanolic extract of *A. biennis* might partly be related to the high death rate of PC12 cells.

Keywords: *Artemisia biennis*, luteolin, kaempferol, apigenin, PC12 cell line

Introduction

Artemisia biennis Willd. (Compositae) is one of 34 *Artemisia* species growing wild in Iran.¹ *Artemisia* spp are used in Iranian folk medicine as eupeptic, emmenagogue, antispasmodic, haemostatic and laxative.^{2,3} Previous phytochemical investigations on *A. biennis* focused on volatile constituents which led to identification of camphor and (E)-beta-farnesene as the major components of essential oils obtained from Iranian and Canadian habitats, respectively.^{4,5} Dichloromethane and ethanolic fractions of *A.biennis* were reported to have cytotoxicity and

leishmanicidal activities, respectively⁶⁻⁸ while no sesquiterpene lactone has been detected in the terpenoid extract of the species.⁹ Another studies on different extracts of *A. biennis* showed the superiority of hydroethanolic and ethyl acetate extracts in antioxidant and antimalarial cell free assays, respectively.^{10, 11} In another research, antioxidant capacity was evaluated in seven fractions (A-G) of defatted hydroethanolic extract of *A. biennis* aerial parts via cell-based methods (in the rat pheochromocytoma cell line PC12). The results indicated that particularly a fraction named F (which was eluted by 80% methanol in water in a reversed phase vacuum liquid chromatography (VLC) system,), possessed notable inhibitory effect on viability of PC12 cell line.¹²

Here, we report the isolation and identification of three flavonoids (1 – 3) from this fraction of the defatted hydroethanolic extract of *A. biennis* aerial parts.

MATERIALS AND METHODS:

Chemicals

LiChroprep[®] RP-18 (15-25 μm) was purchased from Merck and all the solvents used for extraction, fractionation and purification from Scharlau. Deuterated methanol was obtained from Armar AG.

General experimental procedures

HPLC was carried out on a Young Lin apparatus equipped with a binary pump (YL 911S) and a PDA detector (YL 9160) using Vertica (RP₁₈, 250×30 mm i.d., 10 μm) column. A Supelco (RP₁₈, 250×10 mm i.d., 5 μm) column was replaced for final purification. Electron Impact Mass Spectrometry (EIMS) were performed on a 5973 Network Mass Selective Detector (Agilent

technology). The NMR spectra were recorded on a Bruker Avance AV 400, using CD₃OD as the solvent and residual solvent signal used as internal standard.

Plant material

Aerial parts of *Artemisia biennis* Willd. were collected from Zoshk (Razavi Khorasan, Iran) in September 2010. Sample was identified by Dr V. Mozaffarian (Research Institute of Forest and Rangelands, Tehran, Iran). The voucher specimen (No. 12570) has been deposited in the herbarium, Department of Pharmacognosy, Faculty of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran.

Extraction and fractionation

In this study, the fraction residues obtained in the previous studies ^{10, 12} were used. Briefly, hydroethanolic extract was obtained by maceration after successive extraction of dried and ground aerial parts with petroleum ether (40-60), dichloromethane, ethyl acetate, ethanol, and equal mixture of water-ethanol. The fractionation of hydroethanolic extract was carried out by reversed phase VLC method with rising ratios of methanol in water as a mobile phase. A fraction (F) which was eluted by 80% methanol in water was used for further phytochemical study.

Isolation procedure

Primary purification of the fraction F (660 mg) by semi-preparative HPLC (mobile phase: 0-5 min, 50 % MeOH in (H₂O, 0.1 % formic acid), 5-45 min, MeOH from 50 to 90% in (H₂O, 0.1 % formic acid), 45 -48 min, 90% MeOH in (H₂O, 0.1 % formic acid), 48 -49 min MeOH from 90 to 100% in (H₂O, 0.1 % formic acid), 49-55 min 100% MeOH, flow rate 8 ml/min, detection at 220, 254 and 360 nm) yielded eight subfractions F1-F8 (Retention times= 35.0, 37.5, 41.2, 42.3, 43.5, 50.1, 51.0 and 52.1 min, respectively). Further purification of F2 (13.3 mg) by semi-

preparative HPLC (mobile phase: 0–30 min, MeOH from 40 to 60 % in (H₂O, 0.1 % formic acid), 30–31 min MeOH from 60 to 100% in (H₂O, 0.1 % formic acid), 31–37 min 100% MeOH, flow rate 3 ml/min, detection at 220, 254 and 360 nm) yielded compound 1 (0.9 mg; t_R = 26.6 min). Final purification of the F3 (41.0 mg) by semi-preparative HPLC (mobile phase: 0–22.5 min, MeOH from 50 to 65 % in (H₂O, 0.1 % formic acid), 22.5–23 min MeOH from 65 to 100% in (H₂O, 0.1 % formic acid), 23–30 min 100% MeOH, flow rate 3 ml/min, detection at 220, 254 and 360 nm) afforded an additional amount of compound 1 (8.0 mg; t_R = 15.6 min) and compound 2 (1.2 mg; t_R = 15.8 min). The same procedure was followed to isolate and purify compound 3 (23.5 mg; t_R = 20.3 min) from F4 (140.2 mg) (Figure 1). Structure elucidation of isolated compounds was achieved using spectroscopic techniques including EIMS and proton NMR experiments.




Fig 1. Chromatograms of (A) primary purification of the fraction F, (B) purification of the compound 1, (C) purification of the compound 2 and an additional amount of the compound 1, and (D) purification of the compound 3

Results and Discussion

Using combination of solid phase extraction (SPE) and HPLC methods, three flavonoids were isolated from the defatted hydroethanolic extract of *A. biennis* and their structures were elucidated by means of spectroscopic analysis including EIMS and ^1H -NMR. All spectroscopic data were in agreement with those given in the literature.¹³⁻¹⁵

Compounds 1 to 3 were defined to be luteolin, kaempferol and apigenin, respectively (Figure 2).




Fig. 2. Chemical structures of isolated flavonoids from the aerial parts of *A. biennis*

Compound 1

Luteolin (=2-(3,4-dihydroxyphenyl)- 5,7-dihydroxy-4-chromenone); yellow crystalline solid, ^1H -NMR (CD_3OD , 400 MHz): 6.22 (1H, d, $J = 2.0$ Hz, H-6), 6.45 (1H, d, $J = 2.0$ Hz, H-8), 6.55 (1H, s, H-3), 6.92 (1H, d, $J = 8.8$ Hz, H-5'), 7.39 (1H, d, $J = 2.0$ Hz, H-2'), 7.40 (1H, dd, $J = 2.0, 8.8$ Hz H-6').

EIMS: m/z 286 $[\text{M}]^+$.

Compound 2

Kaempferol (=3,5,7-trihydroxy-2-(4-hydroxyphenyl)-4*H*-chromen-4-one); yellow crystalline solid, ¹H-NMR (CD₃OD, 400 MHz): 6.20 (1H, d, *J* = 2.0 Hz, H-6), 6.41 (1H, d, *J* = 2.0 Hz, H-8), 6.93 (2H, d, *J* = 8.4 Hz, H-3',5'), 8.11 (2H, d, *J* = 8.4 Hz, H-2', 6').

EIMS: *m/z* 286 [M]⁺.

Compound 3

Apigenin (=5,7-Dihydroxy-2-(4-hydroxyphenyl)-4*H*-1-benzopyran-4-one); yellow crystalline solid, ¹H-NMR (CD₃OD, 400 MHz): 6.22 (1H, d, *J* = 2.0 Hz, H-6), 6.47 (1H, d, *J* = 2.0 Hz, H-8), 6.61 (1H, s, H-3), 6.95 (2H, d, *J* = 8.8 Hz, H-3', 5'), 7.87 (2H, d, *J* = 8.8 Hz, H-2', 6').

EIMS: *m/z* 270 [M]⁺.

The EIMS molecular ion peak of compounds 1 and 2 observed at *m/z* 286 a.m.u, consistent with the molecular formula of C₁₅H₁₀O₆. In the ¹H-NMR spectrum of compound 1, the presence of a singlet with the chemical shift of 6.55 ppm was indicative of a flavone structure. Substitution patterns of hydroxyl groups in ring A were similar for both compounds while they differed in number of hydroxyl substitution groups on ring B. The EIMS molecular ion peak of compound 3 observed at *m/z* 270 a.m.u, suggesting the molecular formula of C₁₅H₁₀O₅. In the ¹H-NMR spectrum of compound 3, the presence of a singlet with the chemical shift of 6.61 ppm was indicative of a flavone structure. Substitution patterns of hydroxyl groups in rings A and B were similar to those of compound 2.

These facts, along with the comparison of the rest of spectroscopic data with those reported in literature ¹³⁻¹⁵, allowed identification of 1, 2 and 3 as luteolin, kaempferol and apigenin, respectively. There is no previous report on the presence of three flavonoid aglycones in *A. biennis*, but these phytochemicals have been proven to be present in some other species of the genus like *A. sacrorum*, *A. anomala*, *A. vulgaris* and *A. ludoviciana*.¹⁶⁻¹⁹ Luteolin, apigenin and

quercetin are regarded as the most common flavonoids in the tribe Anthemideae. Interestingly, among more than 160 isolated individual flavonoid components in the genus *Artemisia*, one third of them are derivatives of the flavones luteolin and apigenin.²⁰ In the current study, they have been isolated from a fraction of hydroethanolic extract which had the most powerful inhibitory effect on viability of PC12 cell line, based on a previous report.¹² In a study on neuroprotective effects of six phenolic compounds on H₂O₂-induced oxidative damage in PC12 cells, the cell viabilities of all experimental groups increased except the apigenin treated group. Apigenin nearly exhibited no effect on H₂O₂-induced PC12 cells damage.²¹ Survival (%) of undifferentiated PC12 cells preincubated with apigenin prior to exposure to linoleic acid hydroperoxide in MTT assay was reported to be lower than that of negative control. Mean survival percentages of differentiated PC12 cells preincubated with the same condition and determined by the trypan blue exclusion test were the same for apigenin and negative control.²² The viability of the rat cardiac myoblast cell line H9C2 treated with 13.5 µg/ml of apigenin was significantly lower compared with the control.²³ Apigenin induced micronuclei in a dose dependent manner in peripheral human lymphocytes and it may have potential clastogenic activity on human genetic material *in vitro*.²⁴ Clastogenicity of luteolin and apigenin in Chinese hamster lung cell line V79 has been demonstrated as well.²⁵ Time - dependent viability of human peripheral blood lymphocytes treated *in vitro* with the flavonoids apigenin , kaempferol and luteolin administered at different concentrations was studied and it was indicated that 2.86 µg /ml of kaempferol had the highest cytotoxic effect (after 18 h of incubation), followed by luteolin and apigenin. The ability of the flavonoids to cause oxidative damage in lymphocyte DNA was assumed as a reason for these observations.²⁶ Cytotoxicity of isolated flavonoids in the current study has been previously evaluated against normal cell lines including human lung embryonic

fibroblast cell line TIG-1 and human umbilical vein endothelial cell line HUVEC. In comparison with kaempferol, apigenin (similar IC₅₀ values of 29.7 µg/ml in both of the assays) and luteolin (IC₅₀ values of 30.6 and 16.3 µg/ml, respectively) have demonstrated stronger cytotoxicity toward the cell lines.²⁷

Defatted hydroethanolic extract of *A. biennis* which showed potent free radical scavenging effect and metal chelating activity¹⁰ was fractionated and all the fractions were subjected to cell-free and cell-based antioxidant assays.^{10, 12} Fraction F demonstrated disappointing results in metal chelating activity and viability of PC12 cell line.^{10, 12} In the present study, three flavonoids were isolated and identified from this fraction which corroborated previous reports on the biological effects of these compounds and their distribution in the genus *Artemisia*.

Conclusion

The presence of apigenin, luteolin and kaempferol might partly be related to the reported decrease in viability of PC12 cell line after exposure to a fraction of hydroethanolic extract of *A. biennis*. Flavone aglycones seem to be dominant in this fraction.

Ethical Issues

Not applicable.

Conflict of Interest

The authors have no conflicts of interest to declare.

References

- 1- Mozaffarian V. A Dictionary of Iranian Plant Names. Tehran: Farhang Moaser Publishers; 1998. p. 56-8.

- 2- Miraldi E, Ferri S, Mostaghimi V. Botanical drugs and preparations in the traditional medicine of West Azerbaijan (Iran). *J Ethnopharmacol*, 2001;75(2-3):77-87. doi: 10.1016/S0378-8741(00)00381-0
- 3- Ghorbani A. Studies on pharmaceutical ethnobotany in the region of Turkmen Sahra north of Iran (part 1): general results. *J Ethnopharmacol*, 2005;102(1):58-68. doi:10.1016/j.jep.2005.05.035
- 4- Nematollahi F, Rustaiyan A, Larijani K, Nadimi M. Essential oil composition of *Artemisia biennis* Willd. and *Pulicaria undulata* (L.) C.A. Mey., two compositae herbs growing wild in Iran. *J Essential Oil Res.* 2006;18(3):339–41. doi: 10.1080/10412905.2006.9699106
- 5- Lopes-Lutz D, Alviano DS, Alviano CS, Kolodziejczyk PP. Screening of chemical composition, antimicrobial and antioxidant activities of *Artemisia* essential oils. *Phytochemistry*. 2008;69(8):1732-8. doi:10.1016/j.phytochem.2008.02.014.
- 6- Emami A, Zamani Taghizadeh Rabe SH, Ahi A, Mahmoudi M. Study on toxic effects of *Artemisia* spp. fractions from Iran on human cancer cell lines. *Journal of Zanzan University of Medical Sciences and Health Services*. 2010;18(70):58-67.
- 7- Tayarani-Najaran Z, Makki FS, Alamolhodaie NS, Mojarrab M, Emami SA. Cytotoxic and apoptotic effects of different extracts of *Artemisia biennis* Willd. on K562 and HL-60 cell lines. *Iran J Basic Med Sci*. 2017;20(2):166-71. doi: 10.22038/ijbms.2017.8242.
- 8- Emami SA, Zamani Taghizadeh Rabe S, Ahi A, Mahmoudi M. Inhibitory activity of eleven *Artemisia* species from Iran against *Leishmania* major parasites. *Iran J Basic Med Sci*. 2012;15(2):807-11. doi:10.22038/ijbms.2012.4854

- 9- Iranshahi M, Emami SA, Mahmoud-Soltani M. Detection of Sesquiterpene Lactones in ten *Artemisia* species population of Khorasan provinces. Iran J Basic Med Sci. 2007;10(3):183-8. doi:10.22038/ijbms.2007.5293
- 10- Hatami T, Emami SA, Miraghaee SS, Mojarrab M. Total phenolic contents and antioxidant activities of different extracts and fractions from the aerial parts of *Artemisia biennis* Willd. Iran J Pharm Res. 2014;13(2):551-8. doi:10.22037/ijpr.2014.1518
- 11- Mojarrab M, Naderi R, Heshmati Afshar F. Screening of different extracts from *Artemisia* species for their potential antimalarial activity. Iran J Pharm Res. 2015;14(2):603-8. doi:10.22037/ijpr.2015.1653
- 12- Mojarrab M, Mehrabi M, Ahmadi F, Hosseinzadeh L. Protective effects of fractions from *Artemisia biennis* hydroethanolic extract against doxorubicin- induced oxidative stress and apoptosis in PC12 cells. Iran J Basic Med Sci. 2016;19(5):503-10. doi:10.22038/ijbms.2016.6935
- 13- Markham KR, Geiger, H. ¹H nuclear magnetic resonance spectroscopy of flavonoids and their glycosides in hexadeuterodimethylsulphoxide. In: Harborne JB, editor. The Flavonoids- Advances in Research since 1986. London: Chapman and Hall, 1994. p. 441–73.
- 14- Yassa N, Saeidnia S., Pirouzi R., Akbaripour M., Shafiee A. Three phenolic glycosides and immunological properties of *Achillea millefolium* from Iran, population of Golestan. Daru. 2007;15(1):49-52.
- 15- Hadizadeh F, Khalili N, Hosseinzadeh H, Khair-Aldine R. Kaempferol from Saffron Petals. Iran J Pharm Res. 2003;2(4):251-2. doi:10.22037/ijpr.2010.66

- 16- Yuan H, Lu X, Ma Q, Li D, Xu G, Piao G. Flavonoids from *Artemisia sacrorum* Ledeb. and their cytotoxic activities against human cancer cell lines. *Exp Ther Med.* 2016;12(3):1873-8. doi:10.3892/etm.2016.3556
- 17- Zhang L, Li BG, Tian FR, Zhou WM, Zheng XX, Tian JK. Studies on flavonoids of *Artemisia anomala* S. Moore. *Chin Pharm J.* 2010;45(2):104-7.
- 18- Lee SJ, Chung HY, Maier CG, Wood AR, Dixon RA, Mabry TJ. Estrogenic flavonoids from *Artemisia vulgaris* L. *J Agric Food Chem.* 1998;46(8):3325-9. doi:10.1021/jf9801264
- 19- Liu YL, Mabry TJ. Flavonoids from *Artemisia ludoviciana* var. *ludoviciana*. *Phytochemistry.* 1982;21(1):209-14. doi:10.1016/0031-9422(82)80045-9
- 20- Belenovskaja L. *Artemisia*: the flavonoids and their systematic value. In: D.J.N. Hind, H.J. Beentje, editors. *Compositae: Systematics. Proceedings of the International Compositae Conference, Royal Botanic Gardens, Kew, vol. I, chapter 18, 1996. p. 253-9.*
- 21- Hu Q, Wang D, Yu J, Ma G, Pei F, Yang W. Neuroprotective effects of six components from *Flammulina velutipes* on H₂O₂-induced oxidative damage in PC12 cells. *J Funct Foods.* 2017;37:586-93. doi:10.1016/j.jff.2017.07.043
- 22- Sasaki N, Toda T, Kaneko T, Baba N, Matsuo M. Protective effects of flavonoids on the cytotoxicity of linoleic acid hydroperoxide toward rat pheochromocytoma PC12 cells. *Chem Biol Interact.* 2003;145(1):101-16. doi:10.1016/S0009-2797(02)00248-X
- 23- Zhou Z, Zhang Y, Lin L, Zhou J. Apigenin suppresses the apoptosis of H9C2 rat cardiomyocytes subjected to myocardial ischemia-reperfusion injury via upregulation of the PI3K/Akt pathway. *Mol Med Rep.* 2018;18(2):1560-70. doi:10.3892/mmr.2018.9115

- 24- Noel S, Kasinathan M, Rath SK. Evaluation of apigenin using in vitro cytochalasin blocked micronucleus assay. *Toxicol in vitro*. 2006;20(7):1168-72. doi:10.1016/j.tiv.2006.03.007
- 25- Snyder RD, Gillies PJ. Evaluation of the clastogenic, DNA intercalative, and topoisomerase II-interactive properties of bioflavonoids in Chinese hamster V79 cells. *Environ Mol Mutagen*. 2002;40(4):266-76. doi:10.1002/em.10121
- 26- Rusak G, Piantanida I, Mašić L, Kapuralin K, Durgo K, Kopjar N. Spectrophotometric analysis of flavonoid-DNA interactions and DNA damaging/protecting and cytotoxic potential of flavonoids in human peripheral blood lymphocytes. *Chem Biol Interact*. 2010;188(1):181-9. doi: 10.1016/j.cbi.2010.07.008
- 27- Matsuo M, Sasaki N, Saga K, Kaneko T. Cytotoxicity of flavonoids toward cultured normal human cells. *Biol Pharm Bull*. 2005;28(2):253-9. doi: 10.1248/bpb.28.253