

Journal Pre-proofs

Editorial

Michael H. Abraham and his developed parameters: Various applications in medicine, chemistry and biology

Abolghasem Jouyban ^{a,b}, William E. Acree Jr. ^{c,*}

DOI: 10.34172/PS.2022.1

Please cite this article as: Jouyban A. Acree Jr WE. Abraham and his developed parameters: Various applications in medicine, chemistry and biology. Pharm Sci. 2022. doi:10.34172/PS.2022.1

Received Date: 12 January 2022

Accepted Date: 18 January 2022

This is a PDF file of an article which was accepted for publication in Pharmaceutical Sciences. It is assigned to an issue after technical editing, formatting for publication and author proofing.

Michael H. Abraham and his developed parameters: Various applications in medicine, chemistry and biology

Abolghasem Jouyban ^{a,b}, William E. Acree Jr. ^{c,*}

^aPharmaceutical Analysis Research Center and Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran.

^bFaculty of Pharmacy, Near East University, PO BOX: 99138 Nicosia, North Cyprus, Mersin 10, Turkey.

^cDepartment of Chemistry, University of North Texas, Denton, Texas 76203-5070, USA.

*Corresponding author: bill.acree@unt.edu

Prof. Michael H. Abraham obtained his B.Sc. degree in chemistry from the Northern Polytechnic London and did his Ph.D. at the University College London and he was a Reader at the University of Surrey, and then became a research fellow at University College London, where he obtained his D.Sc. degree in 1974 (1).

Prof. Abraham's first paper indexed in Scopus backs to 1957 (2) and dealt with the chromatographic analysis of peroxides. During 1957-1969 his publications mainly focused on chemistry of organometallic compounds, and in 1969, the first paper dealing with solvent effects on the free energies of the reactants and transition states was published in "Journal of the Chemical Society D: Chemical Communications" (3) when he was working as a Reader at the University of Surrey.

In 1981, Michael published his joint first paper with Prof. Taft (from University of California) and Prof. Kamlet (from a military laboratory in Maryland) dealing with the application of solvatochromic parameters in heterocyclic decomposition of *tert*-butyl halides (4). Robert W. Taft was one of the pioneers of linear free energy relationships (LFERs) who separated the substituent's effects into polar, steric and resonance influences (5). A novel method for obtaining scales of hydrogen-bond acidity (α) (6) and hydrogen-bond basicity (β) (7) of the solvents by the solvatochromic comparison was proposed jointly with Mortimer J. Kamlet in 1976. The general π^* scale of the solvent's polarity based on UV-Vis measurements was suggested in collaboration with Kamlet and Abboud (8). Michael added the Hildebrand solubility parameter as another solvent descriptor for correlation of all types of solvent effects (9). Taft and his co-authors including Michael have also realized that the solvatochromic solvent equations could be set up to

investigate the effects of the solutes, when the solvent system remained constant (10). In 1983, they published the first joint paper dealing with the solubility properties of the solutes in polymers and biological media (11). In 1985, solubilities of a range of nonelectrolytes in blood, plasma, brain, lung, liver, kidney, muscle tissue, and human fat were calculated using the developed descriptors (12). Toxic properties of the compounds were investigated using the developed descriptors in 1986 (13). In 1990, the applicability of the developed descriptors for representation of the upper respiratory tract irritation of male mice by airborne chemicals was shown (14).

In 1992, Michael was awarded the first of two Ebert Prizes for his paper entitled “Role of hydrogen bonding in general anesthesia” (15). The Ebert Prize was established in 1873 which is the oldest pharmacy award. It is presented by the American Pharmaceutical Association to the authors of a published paper in Journal of Pharmaceutical Sciences describing novel and original findings with high probability of significantly impacting the pharmaceutical sciences (16). Two years later, the general form of the Abraham solvation model was applied to investigate partitioning of some drugs between blood and brain (17) and also in chromatographic systems (18). In 1995, skin permeation of molecular solutes was described by Abraham solvation model (19). In 2001, human intestinal absorption of molecular drugs was represented using the Abraham solvation parameters calculated using ABSOLV program (20). In 2002, this publication (20) was awarded the second of the two Ebert Prizes that Michael received for his research publications (16).

In 2004, Abraham and Zhao (21) introduced the specific ionic descriptors (J^+ and J^-) for cations and anions which opened a window for application of the solvation parameters for describing human skin permeability (22), human intestinal absorption (23), artificial membrane retention factors (24) and brain permeation (25) for the ionized forms of the solute molecules. Abraham model correlations have also been reported for describing solute transfer of ions and ionic species from water into more than 30 different organic solvents and into select binary aqueous-organic cosolvent mixtures.

The Abraham solvation model has been successfully applied to correlating the gas to room temperature ionic liquid (RTIL) partition coefficients of volatile organic compounds into more than 100 different RTILs of varying polarity and hydrogen-bonding character (26,27). Abraham, Acree and coworkers (28) extended the predictive applicability of the basic model by splitting the six equation coefficients into a separate cation and anion contribution, which when summed will

yield the respective equation coefficient for the given RTIL. At the writing of this editorial coefficients have been reported for 65 individual cations and 20 individual anions. The published cation and anion coefficients allow one to construct Abraham model predictive equations for 1,300 (65 times 20) different RTILs.

The convulsant/anesthetic activity of 48 compounds has been analyzed using an Abraham solvation equation in 2009 and the results revealed that the predicted convulsant pressure is less than that of anesthetic pressure which is in good agreement with the experimental findings for the compounds studied. The difference between the compound pressure to produce convulsions and anesthesia is quite small for many of the 48 compounds, and so minor structural features can be significant (29). An earlier study (30) had examined the effect of anesthetic structure on inhalation anesthesia of 148 varied anesthetic agents based upon measured values of the minimum alveolar concentration.

In 2011, Abraham and Acree proposed a new method to extend the applicability of the Abraham solvent parameters for calculating the solubility in water-ethanol mixtures in which each solvent composition possesses a given set of the solvent parameters (31). The approach was later extended to binary aqueous-methanol (32) and aqueous-isopropyl alcohol (33) mixtures. In his last paper published in *Journal of Solution Chemistry*, water-solvent partition coefficients of fluorescein at an aqueous pH 3.28 for the solvents octan-1-ol, dichloromethane, trichloromethane, 1,2-dichloroethane, toluene, chlorobenzene and bromobenzene were determined and used to compute its descriptors (34).

Michael was among the most influencing scientist in solution chemistry and his developed descriptors have been used for mathematical representation of many chemical, pharmaceutical and biological processes. Even though Michael formally retired many years ago, he continued to publish many important papers. Near the time of his death he was collaborating with Professor William Green (of Massachusetts Institute of Technology) looking to further the applicability of his general solvation parameter model by examining group contribution and machine learning methods as a means to predict solute descriptors for additional organic and medicinal compounds (35). Once calculated the solute descriptors can be used to predict a wide range of different properties using the mathematical correlations published by Professor Abraham and researchers. Several of these correlations have been described above.

It should be noted that the Abraham model solute descriptors are not simply curve-fitting parameters, but rather the descriptors encode valuable chemical information regarding how the given solute molecule interacts with neighboring solvent molecules. Solute descriptors, denoted as E, S, A, B, V and L in Michael's recent publications (19-35), are defined as follows: E denotes to the solute excess molar refractivity in units of $(\text{cm}^3 \text{ mol}^{-1})/10$, S measures the solute dipolarity/polarizability, A and B pertain to the overall or total hydrogen-bond acidity and basicity, V refers to the McGowan volume expressed in units of $(\text{cm}^3 \text{ mol}^{-1})/100$, and L is defined as the logarithm of the solute's gas-to-hexadecane partition coefficient measured at a reference temperature of 298 K. The solute descriptors are described in greater detail elsewhere (37).

The large solute descriptor (36), blood to brain partition coefficient (38) and skin permeation (22) databases that Michael constructed over the years continue to be used by other researchers working in the pharmaceutical and chemical industries. For example, Yamamoto and coworkers (39) recently examined the absorption of pharmaceutical compounds on several laboratory material surfaces using the Abraham model solute descriptors. As noted by the authors unwanted absorption onto test tubes and other laboratory surfaces can adversely affect the accurate determination of drug concentrations in solution, and lead to miscalculated absorption, distribution, metabolism, and excretion (ADME) properties. Li (40) later utilized published Abraham model water to organic solvent correlations in searching for suitable extraction vehicles (blood simulating solvents) to use in testing the safety of blood-contacting medical devices. Several methods (31-33, 41-43) have employed Abraham model solute descriptors as input parameters for estimating the solubility of medicinal compounds in organic mono-solvents and binary solvent mixtures. The derived expressions provide the pharmaceutical community with an *in silico* means to estimate solubility in the absence of experimental values.

As an indication of the impact that Michael's work had, he continues to be listed among the top 2 % of scientist based on a study titled "August 2021 data-update for 'Updated science-wide author databases of standardized citation indicators'" by Jeroen Baas, Kevin Boyack, and John P.A. Ioannidis (44). He was also a professional painter and spent considerable time painting in the studio, landscape painting and also setting up the art exhibitions (45). In Jan 2021 Michael sadly passed away and the researchers of solution chemistry lost one of the most active leaders of the field. This editorial was prepared according to an invitation from the editor of the Pharmaceutical Sciences as the memorial of the first anniversary of his death.

Authors declare no conflict of interests.

Accepted Manuscript

References:

1. Abraham MH. Scales of solute hydrogen-bonding: Their construction and application to physicochemical and biochemical processes. *Chem Soc Rev.* 1993;22(2):73–83.
2. Abraham MH, Davies AG, Llewellyn DR, Thain EM. The chromatographic analysis of organic peroxides. *Anal Chim Acta.* 1957;17(C):499–503.
3. Abraham MH. Solvent effects on the free energies of the reactants and transition states in the menshutkin reaction of trimethylamine with alkyl halides. *J Chem Soc D Chem Commun.* 1969;(22):1307–8.
4. Abraham MH, Taft RW, Kamlet MJ. Linear Solvation Energy Relationships. 15. Heterolytic Decomposition of the tert-Butyl Halides. *J Org Chem.* 1981;46(15):3053–6.
5. Taft RW. Steric effects in organic chemistry. In: Newman MS, editor. New York: Wiley;
6. Taft RW, Kamlet MJ. The Solvatochromic Comparison Method. 2. The α -Scale of Solvent Hydrogen-Bond Donor (HBD) Acidities. *J Am Chem Soc.* 1976;98(10):2886–94.
7. Kamlet MJ, Taft RW. The Solvatochromic Comparison Method. I. The β -Scale Of Solvent Hydrogen-Bond Acceptor (HBA) Basicities. *J Am Chem Soc.* 1976;98(2):377–83.
8. Kamlet MJ, Abboud JL, Taft RW. The Solvatochromic Comparison Method. 6. The π^* Scale of Solvent Polarities1. *J Am Chem Soc.* 1977;99(18):6027–38.
9. Kamlet MJ, Abboud J-LM, Abraham MH, Taft RW. Linear Solvation Energy Relationships. 23. A Comprehensive Collection of the Solvatochromic Parameters, π , α , and β , and Some Methods for Simplifying the Generalized Solvatochromic Equation. *J Org Chem.* 1983;48(17):2877–87.
10. Kamlet MJ, Doherty RM, Abboud JM, Abraham MH, Taft RW. Linear Solvation Energy Relationships: 36. Molecular Properties Governing Solubilities of Organic Nonelectrolytes In Water. *J Pharm Sci.* 1986;75(4):338–49.
11. Abraham MH, Kamlet MJ, Taft RW, Weathersby PK. Solubility Properties in Polymers and Biological Media. 1. Correlations in Various Media of the Solubilities of Gaseous Anesthetics and Other Solutes with a Number of Solute Parameters. *J Am Chem Soc.* 1983;105(23):6797–801.
12. Abraham MH, Kamlet MJ, Taft RW, Doherty RM, Weathersby PK. Solubility Properties in Polymers and Biological Media. 2. The Correlation and Prediction of the Solubilities of Nonelectrolytes in Biological Tissues and Fluids. *J Med Chem.* 1985;28(7):865–70.

13. Kamlet MJ, Doherty RM, Velth GD, Taft RW, Abraham MH. Solubility Properties in Polymers and Biological Media. 7. An Analysis of Toxicant Properties That Influence Inhibition of Bioluminescence in *Photobacterium phosphoreum* (The Microtox Test). *Environ Sci Technol*. 1986;20(7):690–5.
14. Abraham MH, Whiting GS, Alarie Y, Morris JJ, Taylor PJ, Doherty RM, et al. Hydrogen Bonding 12. A New QSAR for Upper Respiratory Tract Irritation by Airborne Chemicals in Mice. *Quant Struct Relationships*. 1990;9(1):6–10.
15. Abraham MH, Lieb WR, Franks NP. Role of hydrogen bonding in general anesthesia. *J Pharm Sci*. 1991;80(8):719–24.
16. No Title. p. <http://www.jpharmsci.org/ebert-prize>.
17. Abraham MH, Chadha HS, Mitchell RC. Hydrogen bonding. 33. Factors that influence the distribution of solutes between blood and brain. *J Pharm Sci*. 1994;83(9):1257–68.
18. Abraham MH, Chadha HS, Leo AJ. Hydrogen bonding. XXXV. Relationship between high-performance liquid chromatography capacity factors and water-octanol partition coefficients. *J Chromatogr A*. 1994;685(2):203–11.
19. Abraham MH, Chadha HS, Mitchell RC. The Factors that Influence Skin Penetration of Solutes. *J Pharm Pharmacol*. 1995;47(1):8–16.
20. Zhao YH, Le J, Abraham MH, Hersey A, Eddershaw PJ, Luscombe CN, et al. Evaluation of human intestinal absorption data and subsequent derivation of a quantitative structure - Activity relationship (QSAR) with the Abraham descriptors. *J Pharm Sci*. 2001;90(6):749–84.
21. Abraham MH, Zhao YH. Determination of solvation descriptors for ionic species: Hydrogen bond acidity and basicity. *J Org Chem*. 2004;69(14):4677–85.
22. Zhang K, Abraham MH, Liu X. An equation for the prediction of human skin permeability of neutral molecules, ions and ionic species. *Int J Pharm*. 2017;521(1–2):259–66.
23. Abraham MH. Human intestinal absorption - Neutral molecules and ionic species. *J Pharm Sci*. 2014;103(7):1956–66.
24. Abraham MH, Acree Jr. WE. Equations for the partition of neutral molecules, ions and ionic species from water to water-ethanol mixtures. *J Solution Chem*. 2012;41(4):730–40.
25. Abraham MH. The permeation of neutral molecules, ions, and ionic species through membranes: Brain permeation as an example. *J Pharm Sci*. 2011;100(5):1690–701.

26. Abraham MH, Acree WE. Comparative analysis of solvation and selectivity in room temperature ionic liquids using the Abraham linear free energy relationship. *Green Chem.* 2006;8(10):906–915.
27. Jiang B, Horton MY, Acree WE, Abraham MH. Ion-specific equation coefficient version of the Abraham model for ionic liquid solvents: determination of coefficients for tributylethylphosphonium, 1-butyl-1-methylmorpholinium, 1-allyl-3-methylimidazolium and octyltriethylammonium cations. *Phys Chem Liq.* 2017;55(3):358–85.
28. Sprunger LM, Proctor A, Acree Jr. WE, Abraham MH. LFER correlations for room temperature ionic liquids: Separation of equation coefficients into individual cation-specific and anion-specific contributions. *Fluid Phase Equilib.* 2008;265(1–2):104–11.
29. Abraham MH, Acree Jr. WE. Prediction of convulsant activity of gases and vapors. *Eur J Med Chem.* 2009;44(2):885–90.
30. Abraham MH, Acree Jr. WE, Mintz C, Payne S. Effect of anesthetic structure on inhalation anesthesia: Implications for the mechanism. *J Pharm Sci.* 2008;97(6):2373–84.
31. Abraham MH, Acree Jr. WE. Partition coefficients and solubilities of compounds in the water-Ethanol solvent system. *J Solution Chem.* 2011;40(7):1279–90.
32. Abraham MH, Acree WE. Equations for the Partition of Neutral Molecules, Ions and Ionic Species from Water to Water–Methanol Mixtures. *J Solution Chem.* 2016;45(6):861–74.
33. Abraham MH, Acree WE, Rafols C, Roses M. Equations for the Correlation and Prediction of Partition Coefficients of Neutral Molecules and Ionic Species in the Water–Isopropanol Solvent System. *J Solution Chem.* 2021;50(4):458–72.
34. Abraham MH, Sánchez-Moreno R. Determination of Water–Solvent Partition Coefficients for Fluorescein: Evaluation of Descriptors for the Lactone Form and Prediction of Properties. *J Solution Chem.* 2021;50(7):1027–35.
35. Chung Y, Vermeire FH, Wu H, Walker PJ, Abraham MH, Green WH. Group contribution and machine learning approaches to predict Abraham solute parameters, solvation free energy, and solvation enthalpy. *ChemRxiv* [Internet]. 2021;1–39. Available from: <https://chemrxiv.org/engage/chemrxiv/article-details/61d33656d6dcc267874e59e0>
36. Ulrich N, Endo S, Brown TN, Watanabe N, Bronner G, Abraham MH, Goss K-U, UFZ-LSER database v 3.2.1 [Internet], Leipzig, Germany, Helmholtz Centre for Environmental Research-UFZ. 2017 [accessed on 12.01.2022]. Available from <http://www.ufz.de/lserd>.

37. Clarke JED, Mallon L. The determination of abraham descriptors and their application to crop protection research. in: Jeschke P, Krämer W, Schirmer U, Witschel M, (Eds.). Modern Methods in Crop Protection Research, Wiley-VCH Verlag GmbH & Co, 2012.
38. Abraham MH, Ibrahim A, Zhao Y, Acree Jr. WE. A Data Base for Partition of VOCs and Drugs from Blood/Plasma/Serum to Brain, and an LFER Analysis of the Data. *J Pharm Sci.* 2006;95(10):2091-100.
39. Yamamoto E, Tominaga N, Kan-no H, Ando D, Miyazaki T, Izutsu K-I. Evaluation of Drug Sorption on Laboratory Materials with Abraham Solvation Parameters of Drugs and its Prevention. *Pharm Res.* 2021;38(12):2167-77.
40. Li J. Evaluation of blood simulating solvents in extractables and leachables testing for chemical characterization of medical devices based on Abraham general solvation model. *J Mol Liq.* 345:116995/1-116995/7.
41. Jouyban A, Rahimpour E, Acree W. Further analysis on solubility measurement and thermodynamic modeling of benzoic acid in monosolvents and binary mixtures. *Pharm Sci.* 2019;25(2):165-70.
42. Jouyban A, Soltanpour S, Soltani S, Chan H, Acree Jr WE. Solubility prediction of drugs in water-cosolvent mixtures using Abraham solvation parameters. *J Pharm Pharm Sci.* 2007;10(3):263-77.
43. Charlton AK, Daniels CR, Acree Jr. WE, Abraham MH. Solubility of Crystalline Nonelectrolyte Solutes in Organic Solvents: Mathematical Correlation of Acetylsalicylic Acid Solubilities with the Abraham General Solvation Model. *J Solution Chem.* 2003;32(12): 1087-102.
44. No Title [Internet]. Available from: <https://elsevier.digitalcommonsdata.com/datasets/btchxktzyw/3>
45. No Title [Internet]. Available from: https://blogs.rsc.org/nj/category/people/page/6/?doing_wp_cron=1640249849.2666280269622802734375 .