

## **Phytochemical analysis and antioxidant activity of eight cultivars of tea (*Camellia sinensis*) and rapid discrimination with FTIR spectroscopy and pattern recognition techniques**

Fatemeh Yousefbeyk, Heshmatollah Ebrahimi-Najafabadi, Sara Dabirian, Salar Salimi<sup>1</sup>, Farshid Baniardalani, Fatemeh Azmian Moghadam, Saeed Ghasemi

**DOI:**[10.34172/PS.2022.27](https://doi.org/10.34172/PS.2022.27)

**Please cite this article as:** Fatemeh Yousefbeyk, Heshmatollah Ebrahimi-Najafabadi, Sara Dabirian, Salar Salimi<sup>1</sup>, Farshid Baniardalani, Fatemeh Azmian Moghadam, Saeed Ghasemi. Phytochemical analysis and antioxidant activity of eight cultivars of tea (*Camellia sinensis*) and rapid discrimination with FTIR spectroscopy and pattern recognition techniques. Pharm Sci. 2022. doi:[10.34172/PS.2022.27](https://doi.org/10.34172/PS.2022.27)

**Received Date:** 2 February 2022

**Accepted Date:** 26 June 2022

This is a PDF file of an article which was accepted for publication in Pharmaceutical Sciences. It is assigned to an issue after technical editing, formatting for publication and author proofing

**Phytochemical analysis and antioxidant activity of eight cultivars of tea (*Camellia sinensis*) and rapid discrimination with FTIR spectroscopy and pattern recognition techniques**

Fatemeh Yousefbeyk<sup>1</sup>, Heshmatollah Ebrahimi-Najafabadi<sup>2</sup>, Sara Dabirian<sup>3</sup>, Salar Salimi<sup>1</sup>, Farshid Baniardalani<sup>1</sup>, Fatemeh Azmian Moghadam<sup>2</sup>, Saeed Ghasemi<sup>2\*</sup>

<sup>1</sup>*Department of Pharmacognosy, School of Pharmacy, Guilan University of Medical Sciences, Rasht, Iran*

<sup>2</sup>*Department of Medicinal Chemistry, School of Pharmacy, Guilan University of Medical Sciences, Rasht, Iran*

<sup>3</sup>*Department of Pharmaceutical Biotechnology, School of Pharmacy, Guilan University of Medical Sciences, Rasht, Iran*

\*Corresponding author: Saeed Ghasemi, Department of Medicinal Chemistry, School of Pharmacy, Guilan University of Medical Sciences complex, Fouman-Saravan highway, Rasht, Iran, Tel: 00981333486470, Fax: 00981333486475, Postal code: 73774-41941. E-mail: ghasemi\_saeed@yahoo.com

## Abstract

**Background:** Tea (*Camellia sinensis* L.O. Kuntze) is one of the most commonly consumed beverages globally, with several beneficial health effects. The composition of leaves is affected by different factors such as climate and tea cultivar.

**Methods:** In this study, the total phenolic, flavonoid, and tannin contents, and antioxidant activities of eight cultivars of tea growing in Iran were determined. The epigallocatechin gallate (EGCG) contents were measured by high-performance liquid chromatography (HPLC) analysis. The Fourier-transform infrared (FT-IR) spectra were used to construct a supervised pattern recognition model using a genetic algorithm-support vector machine (GA-SVM) for the classification of cultivars.

**Results:** The results indicated a wide variation of total phenolic content (178.04 to 316.09 mg gallic acid equivalents (GAE)/g extract). Also, the flavonoid contents ranged from 25.54 to 41.1 mg quercetin equivalents (QE)/g extract. All the samples had close tannin amounts (ranging from 40.32 to 45.90 mg as tannic acid equivalent (TAE)/g extract). There was a significant linear relationship among total phenolic and flavonoid contents and antioxidant properties. The cultivars DN, PK2, and C.Y.9 had the highest content of phenolic and flavonoid content as well as the best antioxidant activity. The EGCG contents were from 2.66 to 4.12%. The highest amount of EGCG was found in cultivars 282, PK2, and C.Y.9. The discriminative region of FT-IR spectra (1350-1650  $\text{cm}^{-1}$ ) were selected using a GA-SVM. This model showed 100% sensitivity and specificity for training and test sets.

**Conclusion:** The leaves phytochemical compositions and antioxidant effect are deeply dependent on the type of tea cultivars. The cultivars PK2 and C.Y.9 can be considered richer sources of polyphenols, especially EGCG. The eight different cultivars can be classified based on chemical components using the recorded FT-IR spectra.

Keywords: *Camellia sinensis*; EGCG; antioxidant; phenolic compound; pattern recognition;

## Introduction

The tea (*Camellia sinensis* L.O. Kuntze), family Theaceae, is one of the most prevalent beverages consumed globally.<sup>1</sup> It grows in tropical and subtropical regions, which provide sufficient rainfall, enough drainage, and somewhat acidic soil.<sup>2</sup> Statistics indicate that between 2005 and 2007, Iran consists of one of the major tea consuming countries in the world, with approximately 1.6 kg per capita tea consumption.<sup>3</sup>

About 34 thousand hectares of lands have been cultured for tea in the North of Iran (Guilan and Mazandaran provinces), which provides a humid temperate climate and adequate yearly precipitation.<sup>3,4</sup> Recently, tea has attracted considerable attention throughout the world because of different health benefits related to its consumption, including antioxidant, anti-arteriosclerotic, anti-obesity, antiallergic, anti-cancer, antimutagenic, anti-apoptotic, neuroprotective, antihyperglycemic, antimicrobial, and anti-inflammatory activities.<sup>5,6</sup> A large body of research demonstrated that overproduction of free radical species could lead to oxidative damage to vital biomolecules, such as DNA, lipids, and proteins. Consequently, they play a crucial role in many diseases such as cancer, heart disease, multiple sclerosis, neurodegenerative disorders (Alzheimer's disease and Parkinson's disease), diabetes mellitus, cardiovascular diseases, and autoimmune diseases.<sup>5,7,8</sup> In general, plant materials that are rich sources of natural antioxidants, especially polyphenols, can cause protective effects against such disorders.<sup>2,7</sup> The tea polyphenols can exert a significant antioxidant effect by scavenging reactive oxygen and nitrogen species as well as acting as metal chelators.<sup>2</sup>

Wide ranges of bioactive compounds like purine alkaloids (in particular, caffeine and theobromine) and polyphenols are responsible for the beneficial effects of tea.<sup>6,9</sup> It has been reported that green tea leaves contain 10–30% tannins or phenolic substances, such as catechins, flavonols, flavanones, phenolic acids, and glycosides.<sup>10,11</sup> The principal flavonols of tea leaves are quercetin, kaempferol, and myricetin, comprising 2-3% of the water-soluble extract materials of tea.<sup>5</sup>

Furthermore, in fresh young leaves of *C. sinensis*, the main catechins (flavan-3-ols) are epigallocatechin gallate (EGCG), epigallocatechin (EGC), epicatechin gallate (ECG), gallic catechin (GC), epicatechin (EC), and catechin.<sup>2,12</sup> Among these compounds, EGCG is considered the predominant one, which takes up around 3-13% of dry tea leaves, and 50-80% of the total catechins in green tea.<sup>9,13</sup> The composition of fresh tea leaves is affected by different factors such as climate, season, tea varieties, and cultivars, horticultural practices, and leaves age.<sup>2,10,14</sup> Therefore, the study of tea cultivars growing in the different regions is one of the global interests in order to determine samples that possess the highest health benefits.<sup>2,12,15-17</sup>

Plant genetic diversity is essential for developing novel and improved cultivars with appropriate characteristics, such as excellent yield potential, and large seeds.<sup>18</sup> Approximately, 2650 accessions of genetic resources from the tea have been recognized in China, Japan, India, Kenya, and other parts of the world.<sup>14,19</sup> Tea germplasms have typically been characterized by biochemical and morphological descriptors. While these descriptors are valuable for tea varietal group identification, they indicate limited levels of inter and intra-varietal polymorphism and they may not account for all the diversity in the species.<sup>20</sup> Numerous previous studies have been developed for the classification of tea cultivars using pattern recognition techniques based on chemical composition data, like gas chromatographic (GC), high-performance liquid chromatography (HPLC), and capillary electrophoresis.<sup>21</sup> However, these methods are costly and complicated to be applied in the routine work.<sup>22</sup> Infrared (IR) spectroscopy methods have been extensively used because it is a rapid and non-destructive method for qualitative and quantitative analyses of many materials.<sup>23</sup> It is extremely fast, chemical-free, and produces no resultant waste. Fourier-transform infrared spectroscopy (FT-IR) combined with attenuated total reflection (ATR) accessory, can be applied for analysing a vast number of samples in a short time without any sample preparation.<sup>24</sup> The advance in chemometric technologies linked with ATR-FT-IR spectroscopy provides an efficient method for determining the physical and chemical properties of products.

Multivariate data analysis and machine learning techniques are excellent strategies for foodstuff discrimination.<sup>25</sup> Cebi et al. used ATR-FTIR to detect sibutramine adulteration in tea and coffee, based on hierarchical clustering and principal component analysis (PCA).<sup>26</sup> Classification of unifloral honey was attempted by ATR-FTIR and random forest.<sup>25</sup> Gori et al. applied a combination of linear discriminant analysis (LDA) and PCA with a moving window algorithm for the classification of grated cheese, based on their ATR-FTIR data.<sup>27</sup>

In the present study, the total phenolic, flavonoid, and tannin contents as well as antioxidant activities of eight different cultivars of tea (*Camellia sinensis*), including PK2, C.Y.9, D.N, D.T.1, 100, 261, 270, and 282, growing in the North of Iran, were determined for the first time. Furthermore, the quantitative analysis of EGCG, as the main catechin in tea leaves, was performed by developing an HPLC method. Finally, the classification of the eight different tea cultivars was performed based on the ATR-FTIR recorded spectra of the extracted samples using chemometric techniques, including partial least squares-discriminant analysis (PLS-DA) and a genetic algorithm-support vector machine (GA-SVM).

## Materials and methods

### *Reagents and chemicals*

Quercetin, tannic acid, EGCG, gallic acid, 2, 2-Diphenyl-1-picrylhydrazyl (DPPH<sup>\*</sup>), Folin–Ciocalteu's phenol reagent, Butylated hydroxy anisole (BHA), and vitamin E were purchased from Sigma-Aldrich chemical company (USA). Sodium bicarbonate, aluminium trichloride, and polyvinylpolypyrrolidone (PVPP) were purchased from Merck, Germany. HPLC-grade acetonitrile and acetic acid were obtained from DUKSAN, Korea. Other used reagents and solvents were of analytical grade. HPLC grade water was obtained from a Millipore Direct-Q purification system.

### *2.2. Plant material and extraction*

According to the reports of Tea Research Centre of Iran (TRCI), Fashalem station, Rasht, Iran, the most common and popular imported tea cultivars, including P.K.2, C.Y.9, D.N, and D.T.1 were chosen. Also, four Iranian genotypes, growing in this center (100, 261, 270, and 282) were chosen to have a comparison among foreign and Iranian cultivars. The youngest two leaves, plus a terminal bud, were hand-plucked from selected eight different tea cultivars, from TRCI, in June, 2019. Samples were all propagated at the TRCI under the same environmental conditions. The tea cultivars differed in their shoot colour and size of the leaves. The voucher specimens were deposited in the herbarium of the school of pharmacy, Guilan University of medical sciences, Rasht, Iran.

The fresh green tea leaves (100 g, each cultivar) were shade dried at room temperature before powdering. The extraction was performed by the method described by Hu et al.<sup>28</sup> The leaves of each cultivar were brewed with ethanol (75%) in a water bath (30°C for 10 min). Then, the extract was filtered (with Whatman No. 1 filter paper) and the remained leaves were extracted by ethanol (35%) at 90°C for 60 min. The extracts obtained from two steps were mixed and dried entirely under the reduced pressure by rotary evaporator.

#### *Determination of total phenolic compounds*

Total phenolic contents (TPC) of extracts were measured by the Folin-Ciocalteu method.<sup>29,30</sup> In the beginning, each extract (1 ml, in the concentration of 1 mg.ml<sup>-1</sup>) was separately mixed with freshly prepared Folin-Ciocalteu reagent (5 ml, diluted ten-fold with distilled water). After 10 min, sodium bicarbonate solution (4 ml, 75 g.l<sup>-1</sup>) was added. The samples were placed in a dark cabinet for 30 min, at room temperature. The absorbance of each sample was measured using a PerkinElmer ultraviolet-visible (UV-Vis) spectrophotometer (LAMBDA 25) at 765 nm. The absorbance of five known concentrations of gallic acid standard solutions (25, 50, 70, 100, and 200 µg.ml<sup>-1</sup>) was measured at the same wavelength, and the calibration curve was plotted. The TPC were expressed as mg gallic acid equivalents (GAE)/ g extract.<sup>8,31</sup>

#### *Determination of total flavonoid content*

Total flavonoid contents (TFC) of extracts were determined, using the technique described by Saeidnia and Gohari.<sup>32</sup> At first, tea extracts (5 ml, 0.5 mg.ml<sup>-1</sup> each) were mixed with 5 ml of aluminium trichloride (AlCl<sub>3</sub>) (2% in methanol) and stand for 10 min. After that, the absorbance of each mixture was measured at 415 nm using a UV/Vis spectrophotometer. Quercetin was used in different concentrations (10, 25, 50, 75, and 100 µg.ml<sup>-1</sup>) to plot the standard curve. Finally, the TFC was expressed as mg quercetin equivalents (QE)/ g extract.<sup>33</sup>

#### *Determination of total tannin content*

The determination of total tannin content (TTC) was performed using PVPP as the insoluble matrix, which binds to tannin phenolics, to measure tannins.<sup>34-36</sup> The test comprised two steps. In the first step, the stock solution of each sample (1 mg.ml<sup>-1</sup>) was prepared in distilled water. Next, 1 ml of each sample and 0.5 ml of Folin-Ciocalteu reagent (1 N) were mixed. Then, 2.5 ml of sodium carbonate (20%) was added, and the mixture was shaken. The absorbance of samples was measured at 725 nm after 40 min. The calibration curve was plotted using different concentrations of tannic acid (0.02, 0.04, 0.06, 0.08, 0.1, 0.15, 0.2 mg.ml<sup>-1</sup>). The amount of total phenols (X), as tannic acid equivalent (TAE) was calculated based on the calibration curve.

In the second step, the tannins were removed from tannin-containing samples by adding PVPP (100 mg PVPP is sufficient to bind 2 mg of total phenols). The samples were vortexed and kept at 4° C for 15 min before being centrifuged (3000 g for 10 min). In this stage, the supernatants were collected. In this supernatant, only simple phenols other than tannins remained. The measurement of the phenolic contents of the supernatants was performed as mentioned in step 1 to determine the contents of non-tannin phenols (Y). Finally, X-Y indicated mg of tannin as TAE/g extracts.<sup>37</sup>

#### *Antioxidant Activity*

The DPPH radical scavenging activities of tea extracts were measured using the method reported by Fakhrabadi et al.<sup>29</sup> Tea leaves extracts were prepared in a series of test tubes. After that, 1 ml of each sample (in different concentrations) was mixed with 2 ml of DPPH solution. After 30 min, the



absorbance was read at 517. The control contained sample (1 ml) and methanol (up to 3 ml), without DPPH. The blank consisted of 1 ml of methanol added to 2 ml of DPPH solution. Furthermore, the antioxidant activities of BHA, vitamin E, and EGCG were tested as the standard antioxidants. The experiments were repeated three times. The inhibition activity was calculated using the following formula.<sup>38</sup>

$$\% \text{Inhibition} = 100 - \left[ \frac{(\text{sample absorption} - \text{control absorption})}{\text{blank absorption}} \right] \times 100$$

The IC<sub>50</sub> values (designate the concentration of the sample (μg.ml<sup>-1</sup>) causing 50% inhibition) were calculated from the graph plotted for inhibition against sample concentration.<sup>31,39,40</sup>

#### *Quantification of EGCG in the tea extracts*

The quantitative analysis of EGCG was carried out by a Waters Alliance e2695 HPLC apparatus. The detector was a Waters 2998 photodiode array (PDA). A C<sub>18</sub> reversed-phase column (Waters Spherisorb: S5 ODS2, 4×250 mm, particle size 5 μm) equipped with a guard column (Waters Spherisorb, S5 ODS2 4.6×10 mm) was used. The Empower 3 software (Waters, USA) was used for data acquisition, peak integration, and calibrations. All the tea samples were dissolved in methanol: distilled water (1:1) (5 mg.ml<sup>-1</sup>). The mobile phases consisted of (A): acetic acid/water (1:99 v/v) and (B): acetonitrile. Elution was performed using the following gradient program: The solvents composition started at 97% A and changed to 0% A during 50 min. A constant flow rate of 1 ml.min<sup>-1</sup> and a temperature of 25 °C were used. The equilibration time was 10 min. The injection volume for all samples was 100 μl. UV absorption spectra were recorded during the HPLC analysis over the wavelength range of 210–400 nm.

The identification was performed by comparing the retention times (R<sub>t</sub>) of each tea sample and their obtained PDA absorbance spectra with the EGCG standard. Different concentrations of EGCG (10, 20, 30, 40, 50, and 60 μg.ml<sup>-1</sup>) were prepared. The EGCG contents were quantified with a 6-point

calibration curve of standard. All data are reported as means  $\pm$  standard deviations of triplicate independent analyses.

#### *Linearity, limit of detection, and limit of quantitation*

Linearity was assessed through the relationship between the concentrations of EGCG and the absorbance found from the PDA detector using the determination coefficient ( $R^2$ ).<sup>41</sup> The calibration lines were plotted by three replicates of each concentration of EGCG (10, 20, 30, 40, 50, and 60  $\mu\text{g} \cdot \text{mL}^{-1}$ ) to recognize the extent of the total variability of the response that could be described by the linear regression model. Limits of detection (LODs) and limits of quantitation (LOQs) were determined using the expressions  $3.3\sigma/S$  and  $10\sigma/S$ , respectively, where  $\sigma$  is the standard deviation of the residues and  $S$  is the slope of the calibration curve.<sup>42</sup>

#### *Classification of tea extracts*

Sample spectra were recorded using ATR-FTIR (Spectrum Two FT-IR spectrometer with ATR accessory (PerkinElmer)) in  $4000\text{--}780\text{ cm}^{-1}$ . Each cultivar was extracted five times, and five FT-IR spectra were recorded for each extract. Spectral baselines were corrected using asymmetric least squares (P.H.C. Eilers, Analytical Chemistry, 75 (2003) 3631). Noise reduction was performed by Savitzky-Golay filter with second-order polynomial and window size of 21. Each cultivar data was randomly divided into training (70%) and test (30%) sets. The training set (144 samples for all cultivars) was used for obtaining the classification model, and a test set (56 samples for all cultivar classes) was used for external validation. For building the classification models, PLS-DA, and SVM with linear kernel function were applied to the training set for the pattern recognition. Monte Carlo cross-validation (100 times, calibration (70%): validation (30%)) was performed for internal assessment of the models. The ratio of misclassified validation set was used for evaluation of the models. The best model was further combined with a GA to reduce the number of variables and to find the informative spectral regions. Finally, the constructed model was assessed by the test set for its

validity. Furthermore, a chance correlation test was also applied to evaluate robustness of the model. All data processing was performed in MATLAB (version 9.8 R2020a, the MathWorks, Natick, MA).

#### *Statistical analyses*

All the experiments were done in triplicates. The results were expressed as mean  $\pm$  standard deviation (SD). Comparison between groups was carried out using One-way ANOVA followed by Tukey *post hoc* test (SPSS software). Differences were considered significant at  $p < 0.05$ . The correlation between antioxidant properties and phytochemical contents were determined by the Pearson procedure, using the SPSS program (Version 22.0. SPSS Inc., Chicago, USA).  $p$ -Values  $< 0.05$  were considered significant.

### **Results and discussion**

#### *Total phenol, total flavonoid, and total tannin contents*

The TPC of tea extracts from eight cultivars were determined by the Folin- Ciocalteu method. The results are shown in Table 1. The analysis of cultivars showed wide variation from  $178.04 \pm 0.06$  mg GAE/g extract (for 100 cultivar) to  $316.09 \pm 0.01$  mg GAE/g extract (for DN cultivar), based on the standard curve of gallic acid ( $y = 0.00096x - 0.0365$ ,  $R^2 = 0.995$ ). The cultivars DN, PK2, and C.Y.9 contained the highest concentrations of phenolic compounds ( $316.09 \pm 0.01$ ,  $313.94 \pm 0.03$ , and  $301.04 \pm 0.04$  mg GAE/g extract, respectively). Statistical analysis showed that the difference in TPC was significant ( $p < 0.05$ ) when compared among tea cultivars. A 1.77-fold difference in TPC was observed between the highest and lowest-ranked cultivars.

Additionally, the total flavonoid content was determined using the standard curve of quercetin ( $y = 0.0187x - 0.0254$ ,  $R^2 = 0.975$ ). The flavonoid contents ranged from  $25.54 \pm 0.17$  to  $41.1 \pm 0.12$  mg QE/g extract. Statistical analysis depicted that the TFC in three cultivars, including DN, PK2, and

C.Y.9, has no significant difference ( $p>0.05$ ). There was a 1.6-fold difference between cultivars with the highest (C.Y.9) and lowest (282) total flavonoid contents.

The TTC were calculated using calibration curve ( $y = 0.0005x - 0.0281$ ,  $R^2 = 0.995$ ). The results indicated that all the samples had very close amounts of tannins (ranged from  $40.32 \pm 0.04$  to  $45.90 \pm 0.03$  mg TAE/ g of extract). The difference in TTC was statistically significant ( $p < 0.05$ ) when compared among tea cultivars, with the exception of 270 and P.K.2 cultivars ( $p > 0.05$ ).

So far, the total phenolic compounds of tea cultivars have been investigated in different studies. Chan *et al.* reported that the amount of total phenols in young tea leaves, growing in Malaysia, was 72.80 mg GAE/g. Also, the DPPH radical scavenging activity was  $30 \mu\text{g} \cdot \text{ml}^{-1}$ .<sup>2</sup> Kerio *et al.* studied the TPC of 10 green and 18 purple leaf-coloured cultivars, growing in Kenya. Their results indicated that the average amount of total phenols ranged between 16.8% and 32%. Their data is inconsistent with the finding of the present study (17.8% -31.6%).<sup>15</sup> Moreover, the determination of TPC in 27 tea cultivars, growing in China revealed a wide range of variation (121.6–223.7 mg of GAE/g extract).<sup>16</sup>

In another study, several cultivars from Sri Lanka were investigated for their TPC. A range from 178 to 251  $\text{mg} \cdot \text{g}^{-1}$  was reported among different samples. The cultivars DN, PK2, and C.Y.9 had 218.62, 238.03, and 205.65  $\text{mg} \cdot \text{g}^{-1}$  total phenols, respectively.<sup>48</sup> Compared to our findings, these three cultivars have a higher amount of phenolic compounds in Iran.

#### *DPPH radical scavenging activity*

Polyphenols in green tea have been described to be potent antioxidant agents. In the recent years, there has been rising interest in using green tea leaves in food industries as a natural antioxidant, to take the place of the synthetic ones, including BHA, and butylated hydroxytoluene (BHT), which are doubted to cause cancer and their usage has been limited in food products.<sup>2,8,49</sup> These phenolic compounds have strong free-radical scavenging activities due to their one-electron reduction potential. They have the ability to act as hydrogen or electron donors.<sup>2</sup>

DPPH radical scavenging assay has been extensively used to measure the radical scavenging power of plant extracts because of its simplicity, affordability, stability, and reproducibility. The DPPH molecule is a stable free radical with an odd electron, which causes an absorption band at about 517 nm and a visible deep violet colour.<sup>50-52</sup> In this method, the DPPH captures an electron from a free radical scavenging antioxidant (hydrogen/electron donor) that leads to a reduction in the intensity of absorption and the solution is decolorized based on the number of electrons accepted.<sup>51,53</sup>

As is presented in Figure 1, the best radical scavenging activity reported from C.Y.9 cultivar ( $IC_{50}= 22.95 \mu\text{g.ml}^{-1}$ ) followed by P.K.2 and D.N cultivars ( $IC_{50}= 25.99$  and  $29.63 \mu\text{g.ml}^{-1}$ ). The C.Y.9 with the highest antioxidant activity showed 2-fold activity of the lowest-ranked one, 270. Furthermore, the  $IC_{50}$  values of EGCG, BHA, and vitamin E were 7.1, 7.77, and  $14.03 \mu\text{g.ml}^{-1}$ , respectively. The antioxidant activity of all cultivars had significant difference ( $p<0.05$ ) compared to vitamin E and BHA. The comparison among other groups is represented in Figure 1.

Furthermore, correlations and regression analysis were performed between total phenolic, flavonoid, and tannin contents and antioxidant activities of tea extracts. Results indicated that DPPH radical scavenging activity had negative correlations and regression with the phenolic, flavonoids, and tannin contents because that the results are expressed as  $IC_{50}$  values.<sup>54</sup>

Interestingly, the concentration of flavonoids exhibited significant correlation with radical scavenging effect ( $r= -0.817$ ,  $R^2=0.668$ ,  $p=0.007$ ). Moreover, the phenolic contents were significantly correlated and influenced the radical scavenging activity in tea extracts (with  $r= -0.646$ ,  $R^2=0.417$ ,  $p=0.042$ ). However, the TTC did not reveal a significant correlation with antioxidant effect ( $r= -0.079$ ,  $R^2= 0.0062$ ,  $p=0.426$ ).

In a study, the  $IC_{50}$  value of green tea extract collected from Japan was reported as  $194.6 \pm 20.5 \mu\text{g.ml}^{-1}$ .<sup>55</sup> Bastos *et al.* investigated the antioxidant activity of green tea, growing in Brazil, and they reported an  $IC_{50}$  of  $88.36 \pm 0.7 \mu\text{g.ml}^{-1}$ .<sup>51</sup> Jain *et al.* reported that the green tea, grown in India, had the

IC<sub>50</sub>= 90.86 µg.ml<sup>-1</sup>, in this essay.<sup>56</sup> Chan *et al.* revealed that the tea leaves growing in Selangor, Malaysia, had an antioxidant activity with an IC<sub>50</sub> of 35 µg.ml<sup>-1</sup>.<sup>2</sup>

#### *Quantitative measurement of EGCG in different tea cultivars*

EGCG, as the most common catechin in green tea leaves, consists of about 50% of the total catechin content and shows the highest antioxidant effect among other catechins. In this study, the quantitative analysis of EGCG content in different tea samples was performed by developing an HPLC method. Figure 2 depicts a typical HPLC chromatogram of tea extracts. The EGCG was detected at R<sub>t</sub>= 26 min. The amount of EGCG ranged from 2.66 to 4.12 %, with an average of 3.06%. As is presented in table 2, the highest concentration of this compound was detected in cultivar 282 (41.24 mg/g extract), followed by cultivar PK2 and C.Y.9 (32 and 30 mg/g extract, respectively). The comparison among different groups is represented in Table 2. No significantly correlative relationships were found between EGCG content and antioxidant activity ( $r=-0.197$ ,  $R^2=0.3067$ ,  $p=0.32$ ).

In another study, Kerio *et al.* reported the amount of EGCG in eleven different cultivars of green leaves tea ranged from 0.46% to 2.92%, with an average of 1.28%.<sup>15</sup> Bazinet *et al.* showed that the tea samples from Quebec, Canada, had 4% EGCG.<sup>57</sup> In another study, Areba *et al.* demonstrated that green tea samples in Kenya had 7% EGCG.<sup>58</sup> The results of our work are in line with the previous studies.

The results of method validation for analysis of EGCG according to linearity, LOD and LOQ indicated that the developed method was valid (Table 3). Outstanding linearity was between 10 and 60 µg.ml<sup>-1</sup>, respectively, with  $R^2=0.994$  for EGCG.

#### *FT-IR spectroscopy*

FT-IR spectra of cultivars were used for the determination of functional groups of biomolecules, existing in tea extracts. The FT-IR spectrum of all tea extracts (Figure 3) showed a broad peak from 3050 to 3600 cm<sup>-1</sup>, which can be assigned to O-H and N-H stretching modes. It can prove the presence of some polyphenol, polysaccharides, and amide groups of proteins. The peaks at 2924 and 2858 cm<sup>-1</sup>

result from asymmetric and symmetric stretching vibrations of aliphatic C-H. Another band at 1688  $\text{cm}^{-1}$  is related to the C=O stretching vibration. The characteristic peak at 1613  $\text{cm}^{-1}$  corresponds to the bending vibration of amide I (N-H) of proteins, stretching vibration of C=C groups, and C-H bending vibration of alkenes. The out-of-plane bending vibration of N-H amide II and III, as well as C-N stretching vibration of proteins appear at 1537 and 1345  $\text{cm}^{-1}$ , respectively. The absorption band at 1448  $\text{cm}^{-1}$  can be ascribed to the bending vibration of  $\text{CH}_2$  alkanes and  $\text{COO}^-$  of carboxylate moiety from proteins and lipids, and rocking vibrations of C-H bonds of cis-disubstituted alkenes. Two peaks at 1233 and 1245  $\text{cm}^{-1}$  are related to the stretching vibration of C-O group of esters. The bands in 1140 and 1205  $\text{cm}^{-1}$  can be attributed to C-H bending and C-H deformation vibrations of fatty acids. The characteristic peak at 1060 shows the stretching vibrations of C-O. Also, all peaks below 1000  $\text{cm}^{-1}$  correspond to the bending modes of  $\text{sp}^2$  C-H of alkenes and aromatic rings.

#### *Pattern recognition*

PLS-DA as a linear classification technique is a combination of the characteristics of PLS regression with the discrimination power of a classification method. This method is based on searching by PLS regression algorithm for latent variables (LVs) with the highest covariance for the dependent variables. Using PLS-DA, the original variables combine linearly as LVs which reflect the relevant sources of data variability. Subsequently, the graphical visualization, a better understanding of the various data patterns as well as their relations with LV scores and loadings are possible. The coefficients of variables are defined as the loadings in the linear combinations. Consequently, they can be explained as the impact of each variable on each LV, while the coordinates of each sample can be interpreted using the score in the LV projection hyperspace.<sup>67</sup> Zau et al. used PLS-DA identification of pesticide residues in tea.<sup>68</sup>

SVM, as a statistical learning method, was introduced by Vapnik.<sup>69</sup> It is an efficient method for prediction and classification of various data sets. GA simulates the process of evolution and the natural genetics based on randomized searches and optimization methods. It is an acceptable method for

classification, prediction, regression estimation, variable selection, and data reduction. Also, a combination of SVM and GA defined as GA-SVM is an efficient tool for pattern recognition and classification.<sup>70-72</sup>

Average error values of 100 times Monte Carlo cross-validation (MCCV) applied on the training set were 0.01 and 0.14 for SVM and PLS-DA, respectively. It should be mentioned that the PLS-DA latent variables and SVM hyperparameters, including box-constraint and kernel scale, were optimized on each iteration of MCCV. As SVM showed a lower error value, it was chosen for further analysis. Table 4 shows the optimized parameters of GA-SVM.

Then, GA-SVM was applied to the training set to select informative regions. Due to the continuity of spectral data, the procedure of selecting well-defined regions is superior to single variables scattered throughout the spectrum. The spectral data were divided into 14 regions. The IR spectra, together with the corresponding 14 regions are presented in Figure 4. The spectral regions between 1660 and 1380 were selected by GA.

Table 5 indicates the accuracy of the established model for train and test sets. The model sensitivity, specificity and accuracy are surprisingly 100%. The model successfully predicts all samples in their own classes. The confusion plot for the training and test sets is presented in Figure 4. The most important area in the spectrum for distinguishing the origin of the samples was the region 1380–1660  $\text{cm}^{-1}$ , which contains mainly the bands of proteins, and unsaturated lipids.

## Conclusion

Tea polyphenols are responsible for multiple and varied health benefits of tea, including antioxidant and chemopreventive effects. This study showed a considerable variation in total phenolic and flavonoid contents, and antioxidant activities among the eight tested cultivars. This data approved that chemical composition of fresh tea leaves is affected by the types of tea cultivars. A significant correlation was detected between antioxidant activity and total phenolic and flavonoid contents. The cultivars DN, PK2, and C.Y.9 had the highest content of phenolic and flavonoid content as well as the



best antioxidant activity. Consequently, these three cultivars can be suggested as better dietary sources of natural antioxidants, mainly polyphenols, to be used as a functional drink or dietary supplements. All evaluated cultivars had a considerable amount of EGCG (especially 282, DN, PK2, and C.Y.9), which make them better candidates for isolation and purification of EGCG, as the main tea catechin, in further investigations. Although the morphological characteristics of eight cultivars are very similar, their classification can be performed based on chemical components, using chemometric methods linked with ATR-FTIR recorded spectra which is a simple, time-saving, and precise method.

### Acknowledgements

This work was supported by Guilan University of Medical Sciences. This study was approved by the Ethical Committee of Guilan University of Medical Sciences (ID: IR. GUMS. REC. 1398. 289).

### Conflict of Interest

The authors confirm that this article content has no conflicts of interest.

### References

1. Jiang H, Engelhardt UH, Thräne C, Maiwald B, Stark J. Determination of flavonol glycosides in green tea, oolong tea and black tea by uhplc compared to hplc. *Food Chem* 2015;183:30-5. doi: 10.1016/j.foodchem.2015.03.024
2. Chan EWC, Lim YY, Chew YL. Antioxidant activity of camellia sinensis leaves and tea from a lowland plantation in malaysia. *Food Chem* 2007;102(4):1214-22. doi: 10.1016/j.foodchem.2006.07.009
3. Ghoochani M, Shekoohiyan S, Yunesian M, Nazmara S, Mahvi AH. Determination of aluminum and zinc in infusion tea cultivated in north of iran. *J Environ Health Sci Eng* 2015;13(1):49. doi: 10.1186/s40201-015-0196-9
4. Shekoohiyan S, Ghoochani M, Mohagheghian A, Mahvi AH, Yunesian M, Nazmara S. Determination of lead, cadmium and arsenic in infusion tea cultivated in north of iran. *J Environ Health Sci Eng (IJEHSE)* 2012;9(1):37. doi: 10.1186/1735-2746-9-37
5. Wang H, Provan GJ, Helliwell K. Tea flavonoids: Their functions, utilisation and analysis. *Trends Food Sci Technol* 2000;11(4-5):152-60. doi: 10.1016/S0924-2244(00)00061-3
6. Azevedo RSA, Teixeira BS, da Silva Sauthier MC, Santana MVA, dos Santos WNL, de Andrade Santana D. Multivariate analysis of the composition of bioactive in tea of the species camellia sinensis. *Food Chem* 2019;273:39-44. doi: 10.1016/j.foodchem.2018.04.030
7. Phaniendra A, Jestadi DB, Periyasamy L. Free radicals: Properties, sources, targets, and their implication in various diseases. *Indian J Clin Biochem* 2015;30(1):11-26. doi: 10.1007/s12291-014-0446-0
8. Ghasemi S, Koohi DE, Emmamzadehashemi MSB, Khamas SS, Moazen M, Hashemi AK, et al. Investigation of phenolic compounds and antioxidant activity of leaves extracts from seventeen cultivars of iranian olive (*olea europaea* l.). *J Food Sci Technol* 2018;55(11):4600-7. doi: 10.1007/s13197-018-3398-1

9. Yang XR, Ye CX, Xu JK, Jiang YM. Simultaneous analysis of purine alkaloids and catechins in camellia sinensis, camellia ptilophylla and camellia assamica var. Kucha by hplc. *Food Chem* 2007;100(3):1132-6. doi: 10.1016/j.foodchem.2005.11.021
10. Perva-Uzunalić A, Škerget M, Knez Ž, Weinreich B, Otto F, Grüner S. Extraction of active ingredients from green tea (camellia sinensis): Extraction efficiency of major catechins and caffeine. *Food Chem* 2006;96(4):597-605. doi: 10.1016/j.foodchem.2005.03.015
11. Farhoosh R, Golmohammadi GA, Khodaparast MH. Antioxidant activity of various extracts of old tea leaves and black tea wastes (camellia sinensis l.). *Food Chem* 2007;100(1):231-6. doi: 10.1016/j.foodchem.2005.09.046
12. Stewart AJ, Mullen W, Crozier A. On-line high-performance liquid chromatography analysis of the antioxidant activity of phenolic compounds in green and black tea. *Mol Nutr Food Res* 2005;49(1):52-60. doi: 10.1002/mnfr.200400064
13. Yu Y, Deng Y, Lu B-m, Liu Y-x, Li J, Bao J-k. Green tea catechins: A fresh flavor to anticancer therapy. *Apoptosis* 2014;19(1):1-18. doi: 10.1007/s10495-013-0908-5
14. Zeng L, Xiao Y, Zhou X, Yu J, Jian G, Li J, et al. Uncovering reasons for differential accumulation of linalool in tea cultivars with different leaf area. *Food Chem* 2021;345:128752. doi: 10.1016/j.foodchem.2020.128752
15. Kerio L, Wachira F, Wanyoko J, Rotich M. Total polyphenols, catechin profiles and antioxidant activity of tea products from purple leaf coloured tea cultivars. *Food Chem* 2013;136(3-4):1405-13. doi: 10.1016/j.foodchem.2012.09.066
16. Zeng L, Luo L, Li H, Liu R. Phytochemical profiles and antioxidant activity of 27 cultivars of tea. *Int J Food Sci Nutr* 2017;68(5):525-37. doi: 10.1080/09637486.2016.1263834
17. Zhao C-N, Tang G-Y, Cao S-Y, Xu X-Y, Gan R-Y, Liu Q, et al. Phenolic profiles and antioxidant activities of 30 tea infusions from green, black, oolong, white, yellow and dark teas. *Antioxidants* 2019;8(7):215. doi: 10.3390/antiox8070215
18. Bhandari H, Bhanu A, Srivastava K, Singh M, Shreya HA. Assessment of genetic diversity in crop plants-an overview. *Adv Plants Agric Res* 2017;7(3):00255. doi: 10.15406/apar.2017.07.00255
19. Chen L, Zhou Z-X. Variations of main quality components of tea genetic resources [camellia sinensis (l.) o. Kuntze] preserved in the china national germplasm tea repository. *Plant Foods Hum Nutr* 2005;60(1):31-5. doi: 10.1007/s11130-005-2540-1
20. Wachira F, Tanaka J, Takeda Y. Genetic variation and differentiation in tea (camellia sinensis) germplasm revealed by rapid and AFLP variation. *J Hort Sci Biotech* 2001;76(5):557-63.
21. Cai J-x, Wang Y-f, Xi X-g, Li H, Wei X-l. Using FTIR spectra and pattern recognition for discrimination of tea varieties. *Int J Biol Macromol* 2015;78:439-46. doi: 10.1016/j.ijbiomac.2015.03.025
22. Palacios-Morillo A, Alcázar Á, de Pablos F, Jurado JM. Differentiation of tea varieties using UV-vis spectra and pattern recognition techniques. *Spectrochim Acta A Mol Biomol Spectrosc* 2013;103:79-83. doi: 10.1016/j.saa.2012.10.052
23. Hazarika AK, Chanda S, Sabhapondit S, Sanyal S, Tamuly P, Tasrin S, et al. Quality assessment of fresh tea leaves by estimating total polyphenols using near infrared spectroscopy. *J Food Sci Technol* 2018;55(12):4867-76. doi: 10.1007/s13197-018-3421-6
24. Li Z, Li B, Zhao Z, Ma W, Li W, Wang J. Rapid identification and quantitative analysis of polycarboxylate superplasticizers using ATR-FTIR spectroscopy combined with chemometric methods. *Math Probl Eng* 2021;2021. doi: 10.1155/2021/6613382
25. Ciulu M, Oertel E, Serra R, Farre R, Spano N, Caredda M, et al. Classification of unifloral honeys from sardinia (italy) by ATR-FTIR spectroscopy and random forest. *Molecules* 2021;26(1):88. doi: 10.3390/molecules26010088
26. Cebi N, Yilmaz MT, Sagdic O. A rapid ATR-FTIR spectroscopic method for detection of sibutramine adulteration in tea and coffee based on hierarchical cluster and principal component analyses. *Food Chem* 2017;229:517-26. doi: 10.1016/j.foodchem.2017.02.072
27. Gori A, Maggio RM, Cerretani L, Nocetti M, Caboni MF. Discrimination of grated cheeses by fourier transform infrared spectroscopy coupled with chemometric techniques. *Int Dairy J* 2012;23(2):115-20. doi: 10.1016/j.idairyj.2011.11.005
28. Hu J, Zhou D, Chen Y. Preparation and antioxidant activity of green tea extract enriched in epigallocatechin (egc) and epigallocatechin gallate (egcg). *J Agric Food Chem* 2009;57(4):1349-53. doi: 10.1021/jf803143n
29. Golfakhrabadi F, Ardekani MRS, Saeidnia S, Yousefbeyk F, Jamalifar H, Ramezani N, et al. Phytochemical analysis, antimicrobial, antioxidant activities and total phenols of ferulago carduchorum in two vegetative stages (flower and fruit). *Pak J Pharm Sci* 2016;29(2):623-8.

30. Yousefbeyk F, Ghasemi S, Evazalipour M, Dabirian S, Schubert C, Hekmatnia S, et al. Phytochemical analysis, antioxidant, antibacterial, and cytotoxic activities of leaves and roots of *rubus hyrcanus* juz. *European Food Research and Technology* 2022;248(1):141-52. doi: 10.1007/s00217-021-03866-z
31. Yousefbeyk F, Vatandoost H, Golfakhrabadi F, Mirzaee Z, Abai MR, Amin G, et al. Antioxidant and larvicidal activity of areal parts of *scrophularia striata* against malaria vector *anopheles stephensi*. *J Arthropod Borne Dis* 2018;12(2):119.
32. Saeidnia S GA. Pharmacognosy and molecular pharmacognosy in practice: A laboratory desk reference of pharmacognosy for researchers and students. Saarbrücken. Germany: Lambert Academic Publishing; 2012.
33. Golfakhrabadi F, Yousefbeyk F, Mirnezami T, Laghaei P, Hajimahmoodi M, Khanavi M. Antioxidant and antiacetylcholinesterase activity of *teucrium hyrcanicum*. *Pharmacogn Res* 2015;7(Suppl 1):S15. doi: 10.4103/0974-8490.157993
34. FOOD ONTI. Quantification of tannins in tree foliage. 2000.
35. Manayi A, Khanavi M, Saiednia S, Azizi E, Mahmoodpour MR, Vafi F, et al. Biological activity and microscopic characterization of *lythrum salicaria* l. *DARU J Pharm Sci* 2013;21(1):61-8. doi: 10.1186/2008-2231-21-61
36. Saeidnia S, Nikan M, Mirnezami T, Gohari AR, Manayi A. Micromorphological characterizations and phytochemicals contents of some *phlomis* species from iran. *Int J Pharm Pharm Sci* 2015;8(1):157-61.
37. Yousefbeyk F, Hemmati G, Gholipour Z, Ghasemi S, Evazalipour M, Schubert C, et al. Phytochemical analysis, antioxidant, cytotoxic, and antimicrobial activities of golden chamomile (*matricaria aurea* (loefl.) schultz bip). *Zeitschrift für Naturforschung C* 2022. doi: doi:10.1515/znc-2021-0269
38. Tofighi Z, Alipour F, Hadavinia H, Abdollahi M, Hadjiakhoondi A, Yassa N. Effective antidiabetic and antioxidant fractions of *otostegia persica* extract and their constituents. *Pharmaceutical biology* 2014;52(8):961-6. doi: 10.3109/13880209.2013.874463
39. Hamidi M, Ghasemi S, Bavafa Bighdilou B, Eghbali Koohi D, Yousefbeyk\* F. Evaluation of antioxidant, antibacterial and cytotoxic activity of methanol extract from leaves and fruits of iranian squirting cucumber (*ecballium elaterium* (l.) a. Rich). *Res J Pharmacog (RJP)* 2020;7(1):23-9. doi: 10.22127/rjp.2019.190800.1509
40. Yousefbeyk F, Dabirian S, Ghanbarzadeh S, Eghbali Koohi D, Yazdizadeh P, Ghasemi S. Green synthesis of silver nanoparticles from *stachys byzantina* k. Koch: Characterization, antioxidant, antibacterial, and cytotoxic activity. *Particulate Science and Technology* 2022;40(2):219-32. doi: 10.1080/02726351.2021.1930302
41. Bozorgi M, Amin G, Kasebzade S, Shekarchi M. Development and validation of a hplc-uv method for determination of proscillaridin a in *drimia maritima*. *Res J Pharmacog (RJP)* 2016;3(3):1-7.
42. Hajimehdipoor H, Amin GR, Adib N, Rastegar H, Shekarchi M. Development of a validated hplc method for the simultaneous determination of flavonoids in *cuscuta chinensis* lam. By ultra-violet detection. *DARU J Pharm Sci* 2012;20(1):1-6. doi: 10.1186/2008-2231-20-57
43. Azmian Moghadam F, Evazalipour M, Ghasemi S, Kefayati H. Design, synthesis, biological evaluation and docking study of novel 4-anilinoquinazolines derivatives as anticancer agents. *Iran J Chem Chem Eng (IJCCCE)* 2020. doi: 10.30492/IJCCCE.2020.132389.4271
44. Moghadam FA, Evazalipour M, Kefayati H, Ghasemi S. 6, 7-disubstituted-4-anilinoquinazoline: Design, synthesis and anticancer activity as a novel series of potent anticancer agents. *Pharm Sci* 2020. doi: 10.34172/PS.2020.72
45. Ghasemi S, Sharifi S, Mojarrad JS. Design, synthesis and biological evaluation of novel piperazinone derivatives as cytotoxic agents. *Adv Pharm Bull* 2020;10(3):423. doi: 10.34172/apb.2020.051
46. Ghasemi S, Davaran S, Sharifi S. Comparison of cytotoxic activity of I778123 as a farnesyltransferase inhibitor and doxorubicin against a549 and ht-29 cell lines. *Advanced pharmaceutical bulletin* 2013;3(1):73. doi: 10.5681/apb.2013.012
47. Yousefbeyk F, Tabaside J, Ostad S, Salehi Sourmaghi M, Amin G. Investigation of chemical composition and cytotoxic activity of aerial parts of *ziziphora clinopodioides* lam. *Research Journal of Pharmacognosy* 2016;3(2):47-51.
48. Kottawa-Arachchi J, Gunasekare M, Ranatunga M, Punyasiri P, Jayasinghe L. Use of biochemical compounds in tea germplasm characterization and its implications in tea breeding in sri lanka. *J Natl Sci Found (JNSF)* 2013;41(4). doi: 10.4038/jnsfsr.v41i4.6252
49. Dufresne CJ, Farnworth ER. A review of latest research findings on the health promotion properties of tea. *J Nutr Biochem* 2001;12(7):404-21. doi: 10.1016/s0955-2863(01)00155-3
50. Hasan SR, Hossain MM, Akter R, Jamila M, Mazumder MEH, Rahman S. Dpph free radical scavenging activity of some bangladeshi medicinal plants. *J Med Plant Res* 2009;3(11):875-9.
51. Bastos DH, Saldanha LA, Catharino RR, Sawaya A, Cunha IB, Carvalho PO, et al. Phenolic antioxidants identified by esi-ms from yerba mate (*ilex paraguariensis*) and green tea (*camelia sinensis*) extracts. *Molecules* 2007;12(3):423-32. doi: 10.3390/12030423

52. Satoh E, Tohyama N, Nishimura M. Comparison of the antioxidant activity of roasted tea with green, oolong, and black teas. *Int J Food Sci Nutr* 2005;56(8):551-9. doi: 10.1080/09637480500398835
53. Locatelli M, Gindro R, Travaglia F, Coisson J-D, Rinaldi M, Arlorio M. Study of the dpph-scavenging activity: Development of a free software for the correct interpretation of data. *Food Chem* 2009;114(3):889-97. doi: 10.1016/j.foodchem.2008.10.035
54. Muniyandi K, George E, Sathyanarayanan S, George BP, Abrahamse H, Thamburaj S, et al. Phenolics, tannins, flavonoids and anthocyanins contents influenced antioxidant and anticancer activities of rubus fruits from western ghats, india. *Food Science and Human Wellness* 2019;8(1):73-81.
55. Katsube T, Tabata H, Ohta Y, Yamasaki Y, Anuurad E, Shiwaku K, et al. Screening for antioxidant activity in edible plant products: Comparison of low-density lipoprotein oxidation assay, dpph radical scavenging assay, and folin-ciocalteu assay. *J Agric Food Chem* 2004;52(8):2391-6. doi: 10.1021/jf035372g
56. Jain DP, Pancholi SS, Patel R. Synergistic antioxidant activity of green tea with some herbs. *J Adv Pharm Technol Res (JAPTR)* 2011;2(3):177. doi: 10.4103/2231-4040.85538
57. Bazinet L, Labbé D, Tremblay A. Production of green tea egc-and egcg-enriched fractions by a two-step extraction procedure. *Sep Purif Technol* 2007;56(1):53-6. doi: 10.1016/j.seppur.2007.01.014
58. Areba GO, Khalid R, Nyaga SN, Thuo JK, Moseti KO, Karori SM, et al. Antioxidant activity and effects of kenyan tea (camellia sinensis) on the liver function and serum biochemistry in male wistar rats. *Int J Basic Clin Pharmacol* 2018;7(8):1471-9doi: 10.18203/2319-2003.ijbcp20183017
59. Yang G-Y, Liao J, Kim K, Yurkow EJ, Yang CS. Inhibition of growth and induction of apoptosis in human cancer cell lines by tea polyphenols. *Carcinogenesis* 1998;19(4):611-6.
60. Kundu T, Dey S, Roy M, Siddiqi M, Bhattacharya R. Induction of apoptosis in human leukemia cells by black tea and its polyphenol theaflavin. *Cancer Lett* 2005;230(1):111-21.
61. Lambert JD, Yang CS. Mechanisms of cancer prevention by tea constituents. *The Journal of nutrition* 2003;133(10):3262S-7S.
62. Yang CS, Maliakal P, Meng X. Inhibition of carcinogenesis by tea. *Annual review of pharmacology and toxicology* 2002;42(1):25-54.
63. Jalilian F, Chahardoli A, Sadrjavadi K, Fattahi A, Shokoohinia Y. Green synthesized silver nanoparticle from allium ampeloprasum aqueous extract: Characterization, antioxidant activities, antibacterial and cytotoxicity effects. *Adv Powder Technol* 2020;31(3):1323-32. doi: 10.1016/j.appt.2020.01.011
64. Fresco P, Borges F, Diniz C, Marques M. New insights on the anticancer properties of dietary polyphenols. *Med Res Rev* 2006;26(6):747-66. doi: 10.1002/med.20060
65. Xu X-Y, Zhao C-N, Cao S-Y, Tang G-Y, Gan R-Y, Li H-B. Effects and mechanisms of tea for the prevention and management of cancers: An updated review. *Crit Rev Food Sci Nutr* 2020;60(10):1693-705. doi: 10.1080/10408398.2019.1588223
66. Christou C, Agapiou A, Kokkinofa R. Use of ftir spectroscopy and chemometrics for the classification of carobs origin. *Journal of Advanced Research* 2018;10:1-8. doi: <https://doi.org/10.1016/j.jare.2017.12.001>
67. Ballabio D, Consonni V. Classification tools in chemistry. Part 1: Linear models. Pls-da. *Anal Methods* 2013;5(16):3790-8. doi: 10.1039/C3AY40582F
68. Zhu J, Sharma AS, Xu J, Xu Y, Jiao T, Ouyang Q, et al. Rapid on-site identification of pesticide residues in tea by one-dimensional convolutional neural network coupled with surface-enhanced raman scattering. *Spectrochim Acta A Mol Biomol Spectrosc* 2021;246:118994. doi: 10.1016/j.saa.2020.118994
69. Vapnik V. The nature of statistical learning theory: Springer science & business media; 2013.
70. Pourbasheer E, Riahi S, Ganjali MR, Norouzi P. Application of genetic algorithm-support vector machine (ga-svm) for prediction of bk-channels activity. *Eur J Med Chem* 2009;44(12):5023-8. doi: 10.1016/j.ejmech.2009.09.006
71. Fernández-Lozano C, Canto C, Gestal M, Andrade-Garda JM, Rabuñal JR, Dorado J, et al. Hybrid model based on genetic algorithms and svm applied to variable selection within fruit juice classification. *Sci World J* 2013;2013. doi: 10.1155/2013/982438
72. Li M, Dai G, Chang T, Shi C, Wei D, Du C, et al. Accurate determination of geographical origin of tea based on terahertz spectroscopy. *Appl Sci* 2017;7(2):172. doi: 10.3390/app7020172

Table 1. Total phenolic, total flavonoid, and total tannin contents of eight different cultivars of *Camellia sinensis*

Sample	Total phenol <sup>1</sup>	Total Flavonoid <sup>2</sup>	Total tannin <sup>3</sup>
<b>100</b>	178.04±0.06 <sup>a</sup>	25.54±0.17 <sup>a</sup>	40.32±0.04 <sup>a</sup>
<b>261</b>	244.7±0.02 <sup>b</sup>	32.32±0.03 <sup>b</sup>	42.70±0.01 <sup>b</sup>
<b>270</b>	228.72±0.01 <sup>c</sup>	23.51±0.11 <sup>c</sup>	45.60±0.04 <sup>c</sup>
<b>282</b>	288.64±0.02 <sup>d</sup>	22.77±0.11 <sup>d</sup>	45.90±0.03 <sup>d</sup>
<b>P.K.2</b>	313.94±0.03 <sup>e</sup>	40.59±0.03 <sup>e</sup>	45.49±0.07 <sup>c</sup>
<b>C.Y.9</b>	301.04±0.04 <sup>f</sup>	41.1±0.12 <sup>e</sup>	43.94±0.01 <sup>e</sup>
<b>D.N</b>	316.09±0.01 <sup>g</sup>	40.69±0.4 <sup>e</sup>	43.38±0.04 <sup>f</sup>
<b>D.T.1</b>	275.46±0.11 <sup>h</sup>	37.92±0.08 <sup>f</sup>	42.38±0.01 <sup>g</sup>

<sup>1</sup> mg of GAE/g extracts; <sup>2</sup> mg of QE/ g of extracts; <sup>3</sup>mg of TAE /g of extracts; the results are expressed as mean ± SD of three independent experiments. Values with no letters in common within each column were significantly different (p < 0.05).

Table 2. The amount of EGCG in eight different cultivars of tea leaves, determining by analytical HPLC

Sample	EGCG <sup>1</sup>
<b>100</b>	30.29±0.12 <sup>a</sup>
<b>261</b>	27.91±0.07 <sup>b</sup>
<b>270</b>	26.68±0.02 <sup>c</sup>
<b>282</b>	41.27±0.02 <sup>d</sup>
<b>P.K.2</b>	32±0.01 <sup>e</sup>
<b>C.Y.9</b>	31.28±0.21 <sup>f</sup>
<b>D.N</b>	30.33±0.15 <sup>g</sup>
<b>D.T.1</b>	27.08±0.12 <sup>c</sup>

<sup>1</sup> mg/g dried extract ± SD, Values with no letters in common within each column were significantly different (p <0.05).

Table 3. Linear range, LOD, and LOQ parameters of EGCG

Compound	Regression equation	R <sup>2</sup>	LOD ( $\mu\text{g}\cdot\text{ml}^{-1}$ )	LOQ ( $\mu\text{g}\cdot\text{ml}^{-1}$ )	Linear range ( $\mu\text{g}\cdot\text{ml}^{-1}$ )
EGCG	614789x-998265	0.994	0.88	2.66	10-60

Note: LOD: Limit of detection; LOQ: limits of quantitation

Table 4. Parameters of genetic algorithm-support vector machine (GA-SVM).

SVM Parameter						
Parameters	C parameter	Kernel Scale		Box Constraint		Kernel Function
SVM	1	0.006		302.3		Linear
GA Parameter						
	Crossover Fraction	Constraint Tolerance	Population	Selection Function	Elite Count	Mutation Function
GA	0.8	10 <sup>-3</sup>	50	Stochastic Uniform	3	Gaussian



Table 5. Classification results of GA-SVM.

	100	261	270	282	P.K.2	C.Y.9	D.N	D.T	Correct	Correct Identification Rate (%)
<b>Train set</b>										
100	18								18	100%
261		18							18	100%
270			18						18	100%
282				18					18	100%
P.K.2					18				18	100%
C.Y.9						18			18	100%
D.N							18		18	100%
D.T								18	18	100%
<b>Test set</b>										
100	7								7	100%
261		7							7	100%
270			7						7	100%
282				7					7	100%
P.K.2					7				7	100%
C.Y.9						7			7	100%
D.N							7		7	100%
D.T								7	7	100%




Figure 1: The IC<sub>50</sub> values of eight tea cultivars, vitamin E, BHA and EGCG in DPPH radical scavenging test. The IC<sub>50</sub> values are expressed as mean  $\pm$  SD of three independent experiments. Different letters above each bar show that the difference between cultivars was significant ( $p < 0.05$ ).




Figure 2: a) HPLC chromatogram of a typical tea extract (10 mg.ml<sup>-1</sup>; injection volume: 100 µl); b) HPLC chromatogram of pure EGCG (60 µg.ml<sup>-1</sup>). UV detection was performed from 210 to 400 nm; the other analytical conditions are designated in the methodology section. Peak identification: EGCG (R<sub>t</sub>=26 min).




Figure 3: ATR-FT-IR spectra of all extracted cultivars. 14 regions are highlighted and labelled accordingly. Inset: magnified the GA selected regions.




Figure 4: Confusion matrix of GA-SVM for a) training set b) test set. (Classes: 1:D.N, 2: 100, 3: 282, 4:PK2, 5: 261, 6: D.T.1, 7: 270, 8: C.Y.9)