



Research Article

Chemical Analysis and Toxicity Screening of *Phlomis olivieri* Benth. and *Phlomis persica* Boiss. Essential Oils

Farideh Mohammadifar¹, Mohammad-Reza Delnavazi², Nagueess Yassa^{2*}¹Faculty of Pharmacy, International Campus, Tehran University of Medical Sciences (IC-TUMS), Tehran, Iran.²Department of Pharmacognosy, Faculty of Pharmacy and Medicinal Plant Research Center, Tehran University of Medical Sciences, Tehran, Iran.

ARTICLE INFO

Article Type:

Original Research

Article History:

Received: 7 April 2015

Accepted: 15 May 2015

Keywords:

*Phlomis olivieri**Phlomis persica*

Essential Oil

Brine Shrimp Lethality Test

 β -Caryophyllene

Germacrene D

ABSTRACT

Background: *Phlomis olivieri* Benth. and *Phlomis persica* Boiss. (Lamiaceae) are two medicinal species endemic to Iran. In the present study, we investigated the chemical compositions and general toxicity potentials of the essential oils obtained from the aerial parts of these two *Phlomis* species. **Method:** The essential oils of the plants were extracted using hydrodistillation method. CG and CG-MS were applied to analyze the chemical constituents of the essential oils and brine shrimp lethality test (BSLT) was used for the evaluation of general toxicity effects of the essential oils. **Results:** A total of 46 compounds were identified in the plants essential oils, among them β -caryophyllene (25.7%) and germacrene D (19.5) in *P. olivieri* and germacrene D (17.2%) and γ -elemene (15.4%) in *P. persica* were the main compounds. The essential oils of *P. olivieri* and *P. persica* exhibited a moderate toxicity activity (LD_{50} : 24.2 ± 0.5 and 41.6 ± 0.8 $\mu\text{g/ml}$, respectively) in brine shrimp lethality test compared to podophyllotoxin (LD_{50} : 2.8 ± 0.3 $\mu\text{g/ml}$). **Conclusion:** The results of present study report essential oils of *P. olivieri* and *P. persica* as two sesquiterpene rich oils with the moderate toxic activity. The chemical constituents of the analyzed essential oils were also found to be different from those reported from other regions of Iran, which may be due to the existence of possible different chemotypes between the populations of these two species.

Introduction

The genus *Phlomis* L. from Lamiaceae family includes about 113 perennial herbs or shrubs distributed in Asia, Europe and Africa.^{1,2} Some of the *Phlomis* species have been reported for their traditional uses as analgesic, diuretic, tonic, anti-diarrheic agents and to treat various conditions such as gastric ulcer, diabetes, inflammation, hemorrhoids and wounds.³ In Flora of Iran, this genus is represented by 19 species, including *Phlomis olivieri* Benth. and *Phlomis persica* Boiss.⁴

Previous *in vivo* and *in vitro* studies demonstrated antioxidant, antimicrobial and antinociceptive activity of *P. olivieri*, as well as antioxidant, cytoprotective, antinociceptive and antidiabetic effects of *P. persica*.⁵⁻¹⁰

A review on the literature revealed that these two species have also subjected to the various phytochemical investigations.¹¹⁻¹⁸ In 2006, Sarkhail *et al.* reported the isolation and structure elucidation of one flavone glycoside; chrysoeriol-7-O- β -D-glucoside and one phenylethanoid glycoside; verbascoside from *P. olivieri* aerial parts, together with chrysoeriol-7-O- β -D-glucoside, chrysoeriol-7-O- β -D-(3"-E-p-coumaroyl) glucoside and one iridoid glycoside; lamiide, from *P.*

persica aerial parts (11). One dimethoxy flavonoid, namely 6,7-dimethoxy-5-hydroxy flavanone has been also reported from the leaves of *P. olivieri*.¹² Furthermore, there are a number of reports related to essential oils composition of these two species from different regions of Iran.¹³⁻¹⁸ In the present study we investigate chemical composition and general toxicity potentials of the essential oils of *P. olivieri* and *P. persica* aerial parts, collected from the Borujen (Chaharmahal-Bakhtiari province) and Nahavand (Hamedan province) regions, respectively. To our knowledge this is the first report on chemical constituents and general toxicity activity of the essential oil of these two species from the mentioned regions in west of Iran.

Material and Methods

Plant material

The flowered aerial parts of *P. olivieri* and *P. persica* were collected in July 2014 from the Gandoman region (Borujen County, Chaharmahal-Bakhtiari province, west of Iran) and Sarabe-Gian region (Nahavand County, Hamedan province, west of Iran), respectively.

*Corresponding Author: Nagueess Yassa, Tel/Fax: (+98) 21 66959101, E-mail: yasa@sina.tums.ac.ir

©2015 The Authors. This is an open access article and applies the Creative Commons Attribution (CC BY-NC) license to the published article. Non-commercial uses of the work are permitted, provided the original work is properly cited.

The plant specimens were verified by botanist M. Aghaahmadi (Isfahan University, Isfahan, Iran).

Essential oil extraction

The air-dried and grounded plants (300 g each) were individually subjected to essential oil extraction for 3.5 h using hydrodistillation method by a Clevenger-type apparatus. The obtained essential oils were then dehydrated over anhydrous sodium sulfate and stored at 4 °C until analysis.

GC and GC-MS analysis

Essential oils of *P. olivieri* and *P. persica* were analyzed using a HP-6890 gas chromatograph (HP-5MS column with 60m × 0.25mm i.d. and 0.25µm film thickness) coupled with a mass detector (HP-6973). The flow rate of carrier gas (Helium) was 1 ml/min. The oven temperature was initiated at 60 °C for 30 min and was then gradually raised at a rate of 5 °C per minute to 250 °C. The injection temperature was 250 °C and the essential oil sample (1 µl) was injected with a split ratio of 1:90. The MS spectra were acquired by electron ionization at 70 eV. The Kovats retention indices (KI) were calculated for all identified compounds using a homologous series of *n*-alkanes injected in equal conditions to the essential oil samples. Identification of the constituents was based on software matching with the Wiley 7n.L online library and also by direct comparison of the retention indices and MS fragmentation patterns with those reported for standard compounds.¹⁹ The essential oils were also analyzed using an HP-6890 gas chromatograph equipped with a FID detector to get relative percentages of the identified compounds. The FID detector temperature was 290 °C and the operation was conducted under the same conditions as described for GC-MS analysis.

Brine shrimp lethality test

General toxicity potentials of the essential oils were evaluated in the brine shrimp lethality test (BSLT) as described by Mojarrab *et al.* with slight modifications.²⁰ Brine shrimps cysts (*Artemia salina* L.) were hatched in sterile artificial seawater (prepared using sea salt 38 g/l and adjusted to pH 9 using Na₂CO₃) under constant aeration for 48 hours at 30 °C. Essential oils (50 mg) were dissolved in DMSO (250 µl) and tween 80 (one drop) and diluted with freshly prepared artificial sea water to obtain solutions with 1000, 700, 500, 300, 100, 10 and 2 µg/ml concentrations in a series of tubes containing about 20 active nauplii in each. The tubes were placed in a water bath at 30 °C for 24 hours under light, and the surviving nauplii were then counted to obtain the concentration causing 50% lethality (LC₅₀ value). Podophyllotoxin, a known cytotoxic aryltetralin lignan, was also applied as positive control. The assay was performed three times and LC₅₀ value was reported as Mean ± SD.

Results and Discussion

Essential oils composition

The hydrodistillation of the aerial parts of *P. olivieri* and *P. persica* yielded 0.2 and 0.1% (V/W) pale yellowish oils, respectively.

Thirty compounds, representing 94.5% of the total oil, were identified as a result of GC and GC-MS analyses of *P. olivieri* essential oil, among them β-caryophyllene (25.7%) germacrene D (19.5), (E)-β-farnesene (9.4) and α-pinene (9.0) were the main compounds (Table 1). The results also indicated that sesquiterpene hydrocarbons (65.3%) were the main group of constituents in *P. olivieri* oil. A review of previous studies on chemical compositions of this species from different regions of Iran revealed that germacrene D with relative percentages of 28.1, 26.4, 66.1 and 48.0-58.0%, in the plants collected from Damavand, Taleghan, Chalus and Kojour regions, respectively and hexahydrofarnesyl acetone (13.3%) in essential oil of the plant collected from Semirrom region have been reported as the main compounds of *P. olivieri* oils (Table 2), whereas in the present study β-caryophyllene (25.7%) was characterized as the main compound of the essential oil of this species aerial parts collected from Borujen region (west of Iran).¹³⁻¹⁷ β-caryophyllene were found at levels of 0-16.0% in previous studies on *P. olivieri* essential oils.¹³⁻¹⁷

This bioactive sesquiterpene (β-caryophyllene) has been also identified as the main compound of some *Phlomis* species, *P. russeliana* (22.6%), *P. chimerae* (31.6%), *P. leucophracta* (20.2%) and *P. linearis* (24.2%), and has been considered for its wide biological activities such as anti-inflammatory, gastric cytoprotection, antispasmodic, local anaesthetic and hepatoprotective effects.²¹⁻²⁸

GC and GC-MS analyses of *P. persica* essential oil resulted in identification of twenty-eight compounds, accounting for 94.1% of the total essential oil. The results demonstrated that the essential oil was rich in sesquiterpene hydrocarbons (36.3%) with germacrene D (17.2%) and γ-elemene (15.4%) as the main compounds followed by bicyclogermacrene (10.8%) and spathulenol (8.8%). As shown in Table 2, there are differences between the main components of the analyzed *P. persica* essential oil and two respective reports on essential oil of this species.¹⁵⁻¹⁸ Sarkhail *et al.* (2004) reported (E)-β-farnesene (21.7%), germacrene D (32.5%), germacrene B (6.1 %) and β-caryophyllene (5.1%) as the main compounds of the essential oil of *P. persica* aerial part collected from Bojnourd region (northeast of Iran).¹⁸ In another study on *P. persica* from Taleghan region (north of Iran), germacrene D (38.2%), bicyclogermacrene (16.3%) and α-pinene (13.3%) were characterized as the main constituents of its aerial part essential oil.¹⁵

Regarding to the wide distribution of *P. olivieri* and *P. persica* in Iran, climatic and geographic factors could be considered as the main responsible of observed differences in essential oil constituents of various populations of these species.²⁹

Table 1. Chemical composition of essential oils obtained from the aerial parts of *P. olivieri* and *P. persica*.

No.	Compounds ^a	KI ^b	Real area (%)	
			<i>P. Olivieri</i>	<i>P. Persica</i>
1	2-Hexenal	847	1.2	-
2	α -Thujene	926	0.3	-
3	α -Pinene	934	9.0	2.1
4	1-Octen-3-ol	976	0.9	-
5	1,2,4-Trimethyl benzene	1022	0.7	-
6	(Z)- β -Ocymene	1024	0.2	-
7	Limonene	1026	1.3	0.9
8	Benzene acetaldehyde	1037	0.6	-
9	α -Terpinolene	1086	0.2	0.7
10	Linalool	1097	2.0	-
11	Nonanal	1103	0.7	-
12	<i>cis</i> -Verbenol	1137	0.4	-
13	Terpinen-4-ol	1174	2.6	-
14	α -Terpineol	1186	0.8	-
15	Undecanal	1207	-	0.5
16	(3Z)-Hexenyl tiglate	1221	-	0.4
17	1,4,4-Trimethyl-6-methylene-cyclohexene	1239	-	0.9
18	α -Cubebene	1248	-	0.5
19	α -Copaene	1375	0.4	-
20	β -Bourbonene	1385	0.9	1.8
21	(E)- β -Damascenone	1386	-	1.2
22	β -Elemene	1389	0.9	2.9
23	β -Caryophyllene	1419	25.7	-
24	γ -Elemene	1422	0.5	15.4
25	Calarene	1436	-	1.5
26	(E)- β -Farnesene	1455	9.4	-
27	Germacrene D	1485	19.5	17.2
28	(E)- β -Ionone	1487	-	1.4
29	β -Selinene	1491	1.4	-
30	Bicyclogermacrene	1500	2.8	10.8
31	α -Muurolene	1503	-	0.3
32	γ -Cadinene	1515	-	0.7
33	δ -Cadinene	1525	0.9	1.5
34	β -Bisabolene	1534	-	1.0
35	Germacrene B	1559	3.5	1.4
36	Virdiflorene	1563	-	0.8
37	Spatulenol	1577	1.5	8.8
38	Alloaromadendrene	1580	-	4.9
39	Caryophyllene oxide	1582	4.2	-
40	1,10-dehydro-isolongifolene	1607	-	1.5
41	Valencene	1621	-	1.6
42	α -Muurolol	1645	-	3.3
43	β -Eudesmol	1649	2.9	-
44	α -Cadinol	1655	1.4	-
44	Hexahydrofarnesyl acetone	1850	1.3	2.7
46	<i>n</i> -Hexadecanoic acid	1965	-	1.6
	Monoterpene hydrocarbons		12.2	11.6
	Oxygenated monoterpenes		5.8	-
	Sesquiterpene hydrocarbons		65.3	67.0
	Oxygenated sesquiterpenes		11.3	11.5
	Nonterpenes		2.9	4.1
	Total identified		97.5	94.1

^a Identified compounds listed in order of elution from HP-5MS column; ^b Relative Kovats retention indices to C₈-C₂₄ *n*-alkanes on HP-5MS column.

Table 2. The results of essential oil analysis of *P. olivieri* and *P. persica* from the previous and present studies.

Plant	Location	Date	Method	Part	Main compounds (%)	Reference
<i>P. olivieri</i>	Semirom (Isfahan)	May 1988	HD ^a	Aerial part	Hexahydrofarnesyl acetone (13.3%) Spathulenol (11.4%) Germacrene D (9.7%) β -Caryophyllene (6.9%)	(13)
	Damavand (Tehran)	July 2001	SD ^b	Aerial part	Germacrene D (28.1%) β -Caryophyllene (16.1%) α -Pinene (11.7%) β -Selinene (10.2%)	(14)
	Taleghan (Tehran)	June 2002	HD	Aerial part	Germacrene D (26.4%) Bicyclogermacrene (12.7%) α -Pinene (7.7%)	(15)
	Chalous (Mazandaran)	May 2003	HD	Aerial part	Germacrene D (66.1%) β -Selinene (5.1%) α -Pinene (4.2%) β -Caryophyllene (4.2%)	(16)
	Kojour (Mazandaran)	May 2005	HD	Leaf	Germacrene D (58%) α -Pinene (10.4%) Bicyclogermacrene (4.4%)	(17)
Flower				Germacrene D (48%) α -Pinene (19.7%) β -Selinene (10.1%)		
Stem				Germacrene D (57%) α -Pinene (8.6%) 2-Pentadecanone (6.5%)		
Root				Germacrene D (35%) α -Pinene (12.6%) Spathulenol (6.6%)		
<i>P. persica</i>	Borujen (Bakhtiari)	July 2014	HD	Aerial part	β -Caryophyllene (25.7%) Germacrene D (19.5%) (E)- β -Farnesene (9.4%) α -Pinene (9.0%)	Present study
	Bojnourd (Khorasan)	June 2002	HD	Aerial part	(E)- β -Farnesene (21.7%) Germacrene D (32.5%) Germacrene B (6.1%) β -Caryophyllene (5.1%)	(18)
	Taleghan (Alburz)	June 2002	HD	Aerial part	Germacrene D (38.2%) Bicyclogermacrene (16.3%) α -Pinene (13.3%)	(15)
	Nahavand (Hamedan)	July 2014	HD	Aerial part	Germacrene D (17.2%) γ -Elemene (15.4%) Bicyclogermacrene (10.8%) Spathulenol (8.8%)	Present study

^a Hydrodistillation, ^b Steam distillation.

Brine shrimp lethality activity

The results of general toxicity assay of the plant essential oils in brine shrimp lethality test (BSL test) were summarized in Table 3. The results indicated that

the essential oils of *P. olivieri* and *P. persica* possessed a moderate toxic activity on *A. salina* larvae with the LD₅₀ values of 24.2 ± 0.5 and 41.6 ± 0.8 µg/ml, respectively, in comparison with podophyllotoxin

(LD₅₀: 2.8 ± 0.3 µg/ml). The main sesquiterpenes of the essential oils such as germacrene D, γ-elemene, bicyclogermacrene, β-caryophyllene and (E)-β-

farnesene etc. may be involved in the toxic effects of the tested essential oils.

Table 3. General toxicity activity of the *P. olivieri* and *P. persica* essential oils in Brine shrimp lethality test.

	<i>P. olivieri</i> essential oil	<i>P. persica</i> essential oil	Podophyllotoxin
Brine shrimp lethality activity (LD ₅₀ , µg/ml)	24.2 ± 0.5	41.6 ± 0.8	2.8 ± 0.3

Conclusion

The results of this study report essential oils of *P. olivieri* and *P. persica* (aerial part) as two sesquiterpene rich essential oils (especially germacrene D and β-caryophyllene) with the moderate toxic activity in BSL test. The observed differences between the chemical constituents of the analyzed essential oil samples and previous reports from other different regions of Iran also propose the existence of possible chemotypes between different populations of these two species.

Conflict of Interest

The authors report no conflicts of interest.

References

- Li MX, Shang XF, Jia ZP, Zhang RX. Phytochemical and biological studies of plants from the genus *Phlomis*. *Chem Biodivers* 2010;7:283-301. doi: 10.1002/cbdv.200800136
- The Plant List. Version 1.1. Published on the Internet; <http://www.theplantlist.org/1.1/browse/A/Lamiaceae/Phlomis/>. (accessed 5 st April 2015).
- Amor IL-B, Boubaker J, Sgaier MB, Skandrani I, Bhourri W, Neffati A, et al. Phytochemistry and biological activities of *Phlomis* species. *J Ethnopharmacol* 2009;125:183-202. doi:10.1016/j.jep.2009.06.022
- Jamzad Z. *Flora of Iran: Lamiaceae*. Tehran: Research Institute of Forests and Rangelands; 2012.
- Morteza-Semnani K, Saeedi M, Shahani S. Antioxidant activity of the methanolic extracts of some species of *Phlomis* and *Stachys* on sunflower oil. *African J Biotech* 2006;5:2428-2432.
- Morteza-Semnani K, Saeedi M, Mahdavi MR, Rahimi F. Antimicrobial Studies on Extracts of Three Species of *Phlomis*. *Pharm Biol* 2006;44:426-429. doi: 10.1080/13880200600798445
- Sarkhail P, Abdollahi M, Shafiee A. Antinociceptive effect of *Phlomis olivieri* Benth., *Phlomis anisodonta* Boiss. and *Phlomis persica* Boiss. total extracts. *Pharmacol Res* 2003;48:263-266. doi:10.1016/S1043-6618(03)00151-8
- Moein S, Moein M, Khoshnoud MJ, Kalanteri T. In vitro antioxidant properties evaluation of 10 Iranian medicinal plants by different methods. *Iran Red Crescent Med J* 2012;14:771. doi: 10.5812/ircmj.1408
- Moein S, Farzami B, Khaghani S, Moein M, Larijani B. Antioxidant properties and prevention of cell cytotoxicity of *Phlomis persica* Boiss. *DARU J Pharm Sci* 2007;15:83-88.
- Sarkhail P, Abdollahi M, Fadayevevan S, Shafiee A, Mohammadirad A, Dehghan G, et al. Effect of *Phlomis persica* on glucose levels and hepatic enzymatic antioxidants in streptozotocin-induced diabetic rats. *Pharmacog Mag* 2010;6:219-224. doi: 10.4103/0973-1296.66940
- Sarkhail P, Monsef-Esfehani HR, Amin G, Surmaghi MHS, Shafiee A. Phytochemical study of *Phlomis olivieri* Benth. and *Phlomis persica* Boiss. *DARU J Pharm Sci* 2006;14:115-121.
- Zardoost MR, Gheibib S. Extraction and identification of flavonoid in *Phlomis olivieri* Benth. *Iran J Org Chem* 2011;3:751-753.
- Ghassemi N, Sajjadi S, Lame M. Volatile constituents of *Phlomis olivieri* Benth. *DARU J Pharm Sci* 2001;9:48-50.
- Mirza M, Nik ZB. Volatile constituents of *Phlomis olivieri* Benth. from Iran. *Flavour Frag J* 2003;18:131-132. doi: 10.1002/ffj.1156
- Khalilzadeh MA, Rustaiyan A, Masoudi S, Tajbakhsh M. Essential oils of *Phlomis persica* Boiss. and *Phlomis olivieri* Benth. from Iran. *J Essent Oil Res* 2005;17:624-625. doi:10.1080/10412905.2005.9699014
- Sarkhail P, Amin G, Shafiee A. Composition of the essential oil of *Phlomis olivieri* Benth. from north of Iran. *DARU J Pharm Sci* 2006;14:71-74.
- Tajbakhsh M, Rineh A, Khalilzadeh MA. Chemical composition of the essential oils from leaves, flowers, stem and root of *Phlomis olivieri* Benth. *J Essent Oil Res* 2007;19:501-503. doi:10.1080/10412905.2007.9699315
- Sarkhail P, Amin G, Shafiee A. Composition of the essential oil of *Phlomis persica* Boiss and *Phlomis chorassanica* Bunge from Iran. *Flavour Frag J* 2004;19:538-540. doi: 10.1002/ffj.1338
- Adams R. *Identification of essential oils by gas chromatography quadrupole mass spectrometry*. Allured Publishing Corporation, Carol Stream, USA. 2001.
- Mojarrab M, Delazar A, Esnaashari S, Heshmati-Afshar F. Chemical composition and general toxicity of essential oils extracted from the aerial parts of *Artemisia armeniaca* Lam. and *A. incana* (L.) Druce growing in Iran. *Res Pharm Sci* 2013;8:65-69. doi:10.1016/j.foodcont.2008.01.001

21. Demirci F, Guven K, Demirci B, Dadandi M, Baser K. Antibacterial activity of two *Phlomis* essential oils against food pathogens. *Food Control* 2008;19:1159-1164. doi:10.1016/j.foodcont.2008.01.001
22. Celik S, Gokturk RS, Flamini G, Cioni PL, Morelli I. Essential oils of *Phlomis leucophracta*, *Phlomis chimerae* and *Phlomis grandiflora* var. *grandiflora* from Turkey. *Biochem Syst Ecol* 2005;33:617-623. doi:10.1016/j.bse.2004.11.010
23. Demirci B, Dadandi MY, Franz G, Başer K. Chemical composition of the essential oil of *Phlomis linearis* Boiss. & Bal., and biological effects on the CAM-assay: a safety evaluation. *Z Naturforsch C* 2002;58:826-829. doi: 10.1515/znc-2003-11-1214
24. Fernandes ES, Passos GF, Medeiros R, da Cunha FM, Ferreira J, Campos MM, *et al.* Anti-inflammatory effects of compounds alpha-humulene and (-)-trans-caryophyllene isolated from the essential oil of *Cordia verbenacea*. *Europ J Pharmacol* 2007;569:228-236. doi:10.1016/j.ejphar.2007.04.059
25. Tambe Y, Tsujiuchi H, Honda G, Ikeshiro Y, Tanaka S. Gastric cytoprotection of the non-steroidal anti-inflammatory sesquiterpene, beta-caryophyllene. *Planta med* 1996;62:469-470. doi: 10.1055/s-2006-957942
26. Leonhardt V, Leal-Cardoso JH, Lahlou S, Albuquerque AA, Porto RS, Celedônio NR, *et al.* Antispasmodic effects of essential oil of *Pterodon polygalaeflorus* and its main constituent β -caryophyllene on rat isolated ileum. *Fundam Clin Pharm* 2010;24:749-758. doi: 10.1111/j.1472-8206.2009.00800.x
27. Ghelardini C, Galeotti N, Mannelli LDC, Mazzanti G, Bartolini A. Local anaesthetic activity of β -caryophyllene. *II Farmaco* 2001;56:387-389. doi:10.1016/S0014-827X(01)01092-8
28. Calleja MA, Vieites JM, Montero-Meterdez T, Torres MI, Faus MJ, Gil A, *et al.* The antioxidant effect of β -caryophyllene protects rat liver from carbon tetrachloride-induced fibrosis by inhibiting hepatic stellate cell activation. *British J Nutr* 2013;109:394-401. doi: 10.1017/S0007114512001298
29. Buchbauer G. *Handbook of essential oil: science, technology, and applications*. New York, USA: CRC press; 2010.